The Components of Physical Fitness: A Roundtable Statement by the American College of Sports Medicine

DEBORAH RIEBE[1], MEIR MAGAL[2], LEONARD A. KAMINSKY[3], ROBERT ROSS[4], STUART M. PHILLIPS[5], SOFIYA ALHASSAN[6], ROSS ARENA[7], DAVID G. BEHM[8], SARAH M. CAMHI[9], TIMOTHY S. CHURCH[10], ELLEN M. EVANS[11], YURI FEITO[12], AVERY D. FAIGENBAUM[13], STEVEN B. HEYMSFIELD[14], CHARLES H. HILLMAN[15], CATHERINE M. JANKOWSKI[16], ANDREAS KONRAD[17], GARY LIGUORI[18], LEAH E. ROBINSON[19], R. ANDREW SHANELY[20], and CAROL EWING GARBER[21]

[1]Department of Kinesiology, University of Rhode Island, Kingston, RI; [2]Department of Exercise Science, North Carolina Wesleyan University, Rocky Mount, NC; [3]School of Kinesiology, Ball State University, Muncie, IN; [4]School of Kinesiology and Health Studies, Queen's University, Kingston, ON, Canada; [5]Department of Kinesiology, McMaster University, Hamilton, ON, Canada; [6]School of Public Health and Health Sciences, University of Massachusetts, Amherst, MA; [7]Department of Physical Therapy, University of Illinois Chicago, Chicago, IL; [8]School of Human Kinetics and Recreation, Memorial University, St. John's, NL, Canada; [9]Department of Kinesiology, University of San Francisco, San Francisco, CA; [10]Wondr Health, Dallas, TX; [11]Department of Kinesiology, Indiana University, Bloomington, IN; [12]Nea Onnim Consultancy, St. Augustine, FL; [13]Department of Kinesiology and Health Sciences, The College of New Jersey, Ewing, NJ; [14]Pennington Biomedical Research Center, Louisianna State University, Baton Rouge, LA; [15]Institute for Cognitive and Brain Health, Northeastern University, Boston, MA; [16]College of Nursing, University of Colorado, Aurora, CO; [17]Institute of Human Movement Science, Sport and Health, University of Graz, Graz, Austria; [18]Department of Kinesiology, Pennsylvania State University Abington College, Abington PA; [19]School of Kinesiology, University of Michigan, Ann Arbor, MI; [20]Department of Kinesiology, Appalachian State University, Boone, NC; and [21]Department of Biobehavioral Sciences, Teachers College, Columbia University, New York, NY.

ABSTRACT

RIEBE D., M. MAGAL, L. A. KAMINSKY, R. ROSS, S. M. PHILLIPS, S. ALHASSAN, R. ARENA, D. G. BEHM, S. M. CAMHI, T. S. CHURCH, E. M. EVANS, Y. FEITO, A. D. FAIGENBAUM, S. B. HEYMSFIELD, C. H. HILLMAN, C. M. JANKOWSKI, A. KONRAD, G. LIGUORI, L. E. ROBINSON, R. A. SHANELY, and C. E. GARBER. The Components of Physical Fitness: A Roundtable Statement by the American College of Sports Medicine. *Med. Sci. Sports Exerc.,* Vol. 58, No. 8, pp. 1827–1850, 2026. The American College of Sports Medicine has included a description of the components of physical fitness (PF) in their publications for 40 yr. Because new scientific evidence has emerged, the American College of Sports Medicine convened a scientific roundtable to reexamine the components of PF. The scientific roundtable agreed upon standardized definitions and an updated evidence-informed model of PF consisting of five interconnected components of PF: 1) cardiorespiratory fitness, 2) muscular fitness, 3) body composition, 4) neuromotor fitness, and 5) flexibility, with muscular fitness, body composition, and neuromotor fitness further separated into subcomponents. These components met four inclusion criteria: they 1) are changeable by exercise, 2) affect the ability to participate in physical activity or exercise, 3) contribute to health, and 4) can be feasibly assessed in professional practice. Additional overriding themes that emerged from the scientific roundtable included the complexity of PF and the interrelated nature of the components. Furthermore, it was agreed that when applied in practice, PF is best addressed in an individualized manner. Therefore, the components of PF should be considered dynamic as the focus on any given component can shift when considering individual needs, goals, health status, and priorities/interests.

Key Words: BODY COMPOSITION, CARDIORESPIRATORY, COMPONENTS, FLEXIBILITY, MUSCULAR FITNESS, NEUROMOTOR

Attaining and maintaining physical fitness (PF) is an important consideration given its critical association with improved physical and mental health, improved quality of life, reduced risks for chronic disease and disability, increased longevity, and improved performance of movement-based pursuits (1–10). Numerous components of PF have been identified, including agility, balance, body composition, cardiorespiratory fitness (CRF), coordination, flexibility, muscular endurance, muscular strength, power, reaction time, and speed

Address for correspondence: Deborah Riebe, Ph.D., Department of Kinesiology, 50 West Independence Way, University of Rhode Island, Kingston, RI 02881; E-mail: debriebe@uri.edu.
Submitted for publication August 2025
Accepted for publication February 2026

0195-9131/2026/588-1827/1850
MEDICINE & SCIENCE IN SPORTS & EXERCISE®
Copyright © 2026 by the American College of Sports Medicine
DOI: 10.1249/MSS.0000000000003997

(9,11–15). The American College of Sports Medicine (ACSM) has included the components of PF into their Guidelines for Exercise Testing and Prescription since 1986. The list of PF components has remained relatively stable since that time, although it was updated in 2009 when "muscular strength and endurance" were separated into the two distinct components of "muscular strength" and "muscular endurance" (12,13,16–22). The components of PF with definitions, as presented in the last five editions of ACSM's Guidelines for Exercise Testing and Prescription, are shown in Table 1 (12,13,16–22).

For more than 40 yr, the components of PF have been dichotomized into health-related and skill-related (also referred to as performance-related) PF (13,14,23). The dichotomized categorization was driven by seminal papers in the 1980s (14,23), reflecting the growing evidence that linked certain PF parameters (e.g., CRF, muscular strength, muscular endurance, body composition, and flexibility) with improvement in a wide range of health outcomes. Health-related PF has served as a blueprint for multiple physical activity (PA) guidelines and is often used within an exercise prescription framework to improve or maintain health and PF (5,9,11,13,15,24). Meanwhile, skill-related PF (e.g., agility, balance, coordination, reaction time, speed, and power) has been more commonly aligned with exercise training for sport performance or motor function (14,15,23). In the years since these papers were published, research has shown that some of the components considered to be skill based are also associated with health (e.g., balance is associated with mortality, physical function, and falling) (9,15,25).

Despite the widespread use of the term PF, there is no ubiquitous agreement on 1) the definition of PF or 2) which components should be included under the PF umbrella (9,11,15,24,26–28). The lack of consensus about the components of PF is apparent, even within the same publication. Several versions of ACSM's Guidelines on Exercise Testing and Prescription list the components of PF, as seen in Table 1, but also refer to neuromotor fitness as a component of fitness within the same text (12,18,19). There are also significant differences among other prominent publications. For example, the 2018 Physical Activity Guidelines for Americans Scientific Advisory Committee Report lists the components of PF as CRF, musculoskeletal fitness, flexibility, balance, and speed (9). Considering the inconsistencies among these and other documents, the lack of uniform definitions, and advances in science, it is appropriate to reexamine the components of PF. Therefore, the ACSM convened a scientific roundtable of experts (referred to as the expert panel for the remainder of the paper) to review and evaluate the scientific evidence concerning the long-standing and widely used components of the PF paradigm. The scientific roundtable consisted of experts vetted and appointed by the ACSM according to its rigorous procedures and represented a broad range of disciplines of ACSM membership (29).

EXPERT PANEL PROCESS

The scientific roundtable was proposed according to the operating codes, policies, and bylaws of ACSM's Scientific Integration and Leadership Committee (29). The proposal included the scientific rationale and scope for the project and the proposed members of the scientific roundtable, selected for their demonstrated expertise in a wide variety of topics relevant to the project. The final composition of the roundtable and project was approved by Scientific Integration and Leadership Committee and ACSM's Board of Trustees.

The expert panel followed procedures consistent with accepted practices for consensus papers (30). The full expert panel participated in a 2-d virtual meeting. Before the meeting, a broad list of components that might fall under the umbrella of PF, gathered from various guidelines and scientific statements, was compiled. Members of the expert panel were charged with completing an independent literature review in their area of expertise and synthesizing the best available evidence. During the meeting, existing definitions of PF and related terms in the literature were examined and discussed. Having achieved consensus on the definition of PF, topic experts presented findings from their review of literature and provided their expert opinions on potential components

TABLE 1. Physical fitness as described in ACSM's Guidelines for Exercise Testing and Prescription, 8th–12th editions.(12,13,16,18,19,21,22)

Health-related physical fitness components
• Cardiorespiratory endurance: the ability of the circulatory and respiratory system to supply oxygen during sustained physical activity
• Body composition: the relative amounts of skeletal muscle, fat, bone, and other elements of the body
• Muscular strength: the ability of the skeletal muscle to exert force
• Muscular endurance: the ability of skeletal muscle to continue to perform activities without undue fatigue
• Flexibility: the range of motion available at a joint
Performance-related physical fitness components
• Agility: the ability to change the position of the body in space with speed and accuracy
• Coordination: the ability to use the senses, such as sight and hearing, together with body parts in performing tasks smoothly and accurately
• Balance: the maintenance of equilibrium while stationary or moving
• Power: the ability or rate at which one can perform work
• Reaction Time: the time elapsed between stimulation and the beginning of the reaction to it
• Speed: the ability to perform a movement within a short period of time

Used with permission (13).

of PF. The committee made preliminary decisions about which components should be further considered for inclusion in the consensus paper using round-robin feedback and discussion (31,32). Each potential component was discussed among the panel; some were eliminated when the expert panel determined there was limited evidence of an independent effect of PA and exercise on the component. The committee recognized that the absence of scientific evidence or practical applications to support the inclusion of a component did not preclude its potential importance or the possibility that it could be reconsidered in the future.

The expert panel unanimously agreed upon a set of standardized criteria to guide the determination of which components and subcomponents should be included in this consensus statement:

- Can regular participation in PA or exercise alter the component?
- Is the component related to the ability to perform PA or exercise?
- Is the component associated with health or health-related outcomes?
- Are there valid and reliable assessments that can be feasibly used in professional practice?

Following the virtual meeting, smaller topic-specific working groups were formed and met independently to further investigate the components that passed the preliminary review. For each component, the evidence was evaluated according to the four criteria, and the working groups presented the rationale supporting the inclusion or exclusion of a component to the full group for final decision-making by consensus. The final step was to translate the recommendations from the panel into a usable model that could be applied to professional practice. During the writing process, the literature was regularly updated to reflect new scientific evidence. Further, as the paper developed, there was continued discussion of where components and subcomponents belonged, and which would be included in the final draft of the paper.

Each member of the expert panel independently reviewed several versions of the paper in its entirety and provided feedback. Following revisions, the final version was unanimously approved. In this consensus statement, we present an updated model of PF along with the scientific rationale for the proposed recommendations.

DEFINITION OF TERMS

An evaluation of PF necessitates standardized definitions of PF and related terms. PF has been defined in many ways, with considerable overlap among the definitions (9,13,23,24,33–38). After an extensive review and discussion of the available definitions and scientific literature, we propose adapting the definition of Caspersen (23) to this simple yet clear definition: PF is a set of interrelated and malleable attributes individuals have or achieve that relates to their ability to perform PA. PA and exercise are related to PF. PA is defined as any bodily movement produced by the contraction of skeletal muscles that results in energy expenditure above the resting level (23,24,33,39). Exercise is defined as a type of PA consisting of planned, structured, and repetitive movements done to improve or maintain one or more components of PF and to achieve a health benefit (23,33). While the outcomes of this consensus paper apply to both PA and exercise, the word exercise is used henceforth in this document because it connotes intentional PA for the purpose of improving health and fitness (9,33).

GUIDING PRINCIPLES

The expert panel adopted the following foundational principles to be applied to this consensus statement:

The findings from this statement are primarily intended for application to adults

The PF model presented here applies to all adults, including older adults and people living with disabilities and health conditions. While many of the concepts included in this paper can apply to children and adolescents, the youth literature was not reviewed. Developmental and physiological differences in children and adolescents should be considered when applying this information to the pediatric population.

PF is an applied concept to support health and fitness goals for adults of all ages, abilities, and health conditions

PF is central to athletic pursuits, but the training methods and aims for peak competitive athletic performance differ from those of individuals who adopt exercise or recreational sports for enjoyment, health, or other reasons. The fitness concepts presented here can apply to highly competitive athletes; however, our scientific review was focused on nonathletes. For that reason, high-level athletes deserve separate consideration and are not meant to be covered within the scope of this statement.

All components of PF are related to health and movement performance, negating the need for a dichotomous categorization

Distinguishing between health- and skill-related components of PF is no longer justified given mounting evidence demonstrating that components previously considered uniquely skill-related are now firmly established markers of numerous health outcomes, and all are related to the performance of exercise (40–43).

The components of PF are interrelated

The complex relationship among the components of PF suggests they may improve health and PF outcomes in an independent, additive, or synergistic manner. The components are closely linked with each other, often making it difficult to consider any given component in isolation. For example, improvements in muscular strength can improve balance (42,44) and body composition (45), underscoring the synergistic effects of resistance exercise on health- (46) and movement-related outcomes. The intersections and overlap between the components of PF reflect the integration of multiple physiological systems during exercise and reflect the principle of specificity-imposed demands (47). In this consensus statement, the components of PF are viewed on a continuum where often there is considerable overlap and interactional effects.

The relative importance of each PF component depends on the individual

The expert panel considers PF a personal attribute that is best understood when considered in the context of the individual's unique circumstances. Being "physically fit" varies across individuals based on biological factors (48,49), and it can be moderated by environmental, sociocultural, and psychological factors that can affect participation in PA (50–53). For example, for an older adult, being "physically fit" may mean having the ability to perform activities of daily living (ADLs) and instrumental activities of daily living, with the goal of maintaining independent living or to engage in desired leisure activities. Other individuals may have the goal of improving PF to attenuate health risks or to pursue lifestyle activities such as hiking or other recreational pursuits. People living with disabilities or chronic health conditions may have specific limitations to exercise, necessitating an individualized approach to PF. Meeting an individual's PF goals warrants consideration of the relative importance of each component of PF to that individual, and how they may be prioritized to reach individualized desired outcomes. Thus, one person may emphasize CRF, while another may prioritize the components of muscular strength or body composition, and these choices may change over time.

The components of PF can be feasibly assessed in practice

For the PF model to have relevance to professional practice, the components must be able to be feasibly assessed in most health and fitness settings. Therefore, only components that can be assessed with equipment and tests that are commonly available are included in the consensus statement. The assessments cited in this paper serve as examples that can be used in practice; it is not an exhaustive list, nor is it an endorsement of specific assessments.

AN UPDATED MODEL OF PHYSICAL FITNESS

In this section of the consensus statement, an updated model of PF, as it relates to health and fitness, is introduced. The components of PF model consist of five broad categories that meet the standardized inclusion criteria described earlier: 1) CRF, 2) muscular fitness, 3) body composition, 4) neuromotor fitness, and 5) flexibility, with muscular fitness, body composition, and neuromotor fitness further divided into subcomponents. Each component and subcomponent is defined in Table 2. Although multiple definitions for each component were available in the literature, the one presented effectively communicates the concept that is consistent with the four inclusion criteria established by the expert panel. A brief rationale for why each component was included in the updated PF model, based on the standardized criteria outlined earlier in the article, is presented.

Cardiorespiratory Fitness

CRF (Table 2) represents the functional capacity of an individual and is an accepted indicator of overall health (8,72). CRF reflects the integrated function of several bodily systems (e.g., cardiovascular, respiratory, and muscular) involved in the transport, uptake, and utilization of oxygen required to support physical exertion (8,73).

Exercise-induced changes. Systematic reviews and meta-analyses have consistently demonstrated a significant improvement in CRF following structured aerobic exercise interventions in adults with or without disease (74–78).

Ability to perform exercise. CRF is directly related to the ability to perform exercise (79,80). CRF must be sufficiently high to support the wide range of energy demands of daily activities, including ADLs/IADLs, occupation, transportation, and exercise in which an individual participates without excessive fatigue (81–85). Further, low CRF has been linked to poor lower extremity muscular strength (86).

Health-related benefits. Clear, consistent, and convincing evidence shows CRF is inversely associated with all-cause and disease-specific morbidity and mortality (4,87–94). A recent meta-analysis found that for every metabolic equivalent (MET) increase in CRF, there was a 14% and 16% decrease in all-cause and cardiovascular disease mortality, respectively (87). Clinically, low CRF is used as a prognostic indicator of surgical risks, mortality, and functional limitations (95–97). Low CRF also is associated with increased risk of falling (92), poorer lower extremity strength (86), reduced cognitive function (98), higher risk of dementia (98), and increased risk of

TABLE 2. Recommended standardized definitions of the components and subcomponents of physical fitness.

Cardiorespiratory fitness: the capacity of the circulatory and respiratory systems to deliver oxygen to the skeletal muscle mitochondria to meet the energy demands of sustained physical activity without undue fatigue (8,23).

Muscular fitness: a multidimensional construct composed of muscular strength, muscular endurance and muscular power that represents the structure, function and quality of skeletal muscles (54,55).

- Muscular strength: the ability of skeletal muscle to exert maximal force on an external object or resistance under a particular set of biomechanical conditions (54,55).
- Muscular endurance: the ability of skeletal muscle to complete repeated contractions against a submaximal resistance within a set time limit or to fatigue (56,57).
- Muscular power: the ability of skeletal muscle to exert force at a given velocity of movement; the product of force and velocity over a unit of time (58).

Body composition: the relative proportions of the total body mass divided into the two components of fat mass and fat-free mass (59,60,61).

- Fat mass (FM): the mass of body fat and other lipids in the body (60).
- Fat-free mass (FFM): the mass of all nonfat components of the body wherever they occur, including skeletal muscles, bones, organs, ligaments, tendons, and bodily fluids (60).

Neuromotor fitness: the ability of an individual to effectively execute a variety of motor tasks during physical activity that may include a combination of gross and fine motor skills of varying complexity (11,62,63,64).

- Agility: the ability to execute a change of speed or direction through a rapid movement of the body in response to a stimulus (65).
- Balance: the ability to maintain a specific body posture by maintaining, achieving or restoring the center of gravity within the base of support. Includes static and dynamic balance (66).
  - Static balance: the ability to sustain a specific position while the body is stationary (66).
  - Dynamic balance: the ability to sustain a specific position and orientation of the body while part or all of the body is in motion (66).
- Gait speed: the speed of ambulation performed with or without assistance (67,68).

Flexibility: the ability to move a joint through its maximal unrestricted range of motion without pain (69,70,71).

cardiovascular complications (86). Higher CRF is associated with better mental health, with studies reporting an inverse dose–response relationship between CRF and mental health disorders such as anxiety and depression (99,100).

Assessments. Cardiopulmonary exercise testing with the measurement of oxygen uptake (VO_2) and related variables (e.g., carbon dioxide production, pulmonary ventilation, and heart rate) is considered the gold-standard approach to assessing CRF (73,94,101–103), but it is not widely available in all health and fitness settings. Maximal or submaximal exercise tests or field tests using a variety of exercise modalities can estimate CRF by using one or more variables such as work rate, heart rate, time, and distance obtained during or following exercise (11–13,103–105). Additionally, several nonexercise equations can be used to predict CRF (103,106–109). These indirect methods to estimate CRF are more practical than gold-standard approaches and can be applied in a variety of settings. While indirect estimations of CRF are not optimal, they represent feasible ways to estimate CRF in settings that preclude other forms of assessment, and they are sufficiently sensitive to detect change (12,13,110).

Muscular Fitness

Muscular fitness is a collective term that includes the interrelated subcomponents of muscular strength, muscular endurance, and muscular power (Table 2) (13,111–114). All subcomponents of muscular fitness can be developed with consistent participation in resistance exercise (111,113). The improvements in muscular fitness consequent to training result from an array of physiological changes in the trained skeletal muscle that include neuromuscular adaptations; muscular hypertrophy (skeletal muscle cross-sectional area) (111,115,116); improved skeletal muscle quality (amount of strength and power per unit of muscle mass) (117–119); improved arterial function, blood flow, and blood pressure (120,121); and numerous molecular alterations in the skeletal muscle (122,123). The specific training-induced outcomes depend primarily on program design variables including load, repetitions, sets, selected exercises, velocity, duration of rest, and frequency (111,113,124–132).

Exercise-induced changes. Most well-designed resistance exercise programs that include appropriate variations in program design variables result in improvements in both muscular strength and muscular endurance (115,132–135). Resistance exercise specifically designed to increase muscular power improves the ability to generate higher power output (13,126,130,136,137). Further, a basis for developing muscular power is underpinned to a degree by the development of strength, since an individual cannot possess a high level of power without being relatively strong (115,58).

Ability to perform exercise. Higher levels of muscular fitness contribute to improved exercise performance. Muscular strength and muscular power are strongly correlated with jumping, sprinting, and change of direction, and improve recreational sport performance (138–141) including sports such as golf (142), pickleball (143), rowing (144), and running (145). Resistance training significantly improves physical function, the ability to perform ADLs, and gait speed in overweight and obese adults (146) and older adults (113,147–152).

Health-related benefits. Muscular fitness is directly associated with overall health and reduced risks of cardiovascular disease, cancer, disability, and other chronic conditions (45,153). Enhancing muscular fitness is independently associated with numerous health benefits including a reduced risk of all-cause mortality (154–157), positive changes in cardiometabolic disease biomarkers (158–163), reductions in blood pressure (164), improved mental health (165–172), improved cognitive function (173,174), increased bone density (9,175,176), reduced low back pain (177–179), reduced pain associated with osteoarthritis (9,180), reduced risk

of type 2 diabetes (181–185), and improved insulin sensitivity and glucose control in people with type 2 diabetes (186–188). Resistance exercise improves muscular fitness in people with acute and chronic diseases such as people living with cancer (5), stroke (189), muscular dystrophy (190), and HIV (191,192). In older adults, the unique and well-established benefits of improved muscular strength and power are numerous and can mitigate the effects of sarcopenia and physical function declines, slow the progression of certain chronic diseases (e.g., osteoporosis), improve cognitive function and the performance of ADLs, and reduce the risk of falls (154,193–200).

Assessments. Numerous assessments exist to measure muscular fitness. Dynamic muscular strength can be assessed directly with a one-repetition maximum test for the upper body (e.g., chest press) or lower body (e.g., leg press) using free weights or selectorized machines (13,201,202). One-repetition maximum can also be estimated using prediction equations based on multiple repetitions at submaximal loads (203–211) or by using ratings of perceived exertion (205,212,213). Isometric tests of muscular fitness are less time-consuming and usually less complicated alternatives to repetition maximum tests, and these tests generally have good reliability and sensitivity to changes in strength (214). Common isometric test protocols include those using portable force cell (hydraulic) or electronic (force-sensitive resistor) dynamometers (e.g., handgrip or fixed dynamometer) to measure maximal voluntary contraction or rate of force development for measurement of strength or power, respectively, of hands, fingers, lower extremities, and other parts of the body (12,215–219). There are also functional isometric tests that measure muscular strength, endurance, or power that are appropriate for a wide array of adults with varying functional ability, such as the plank test (prone bridge test), wall sit test, and back strength test, among others (220). Submaximal tests of the subcomponents of muscular fitness include the YMCA bench press (strength and endurance), push-up and pull-up test (upper extremity strength and endurance), stair climb (power), chair stand test (lower extremity strength and power), and 30-s arm curl test (upper extremity strength and power) (210,211,221–225).

Body Composition

Body composition (Table 2) refers to the distribution of the components that constitute the human body. The primary components most often considered in the assessment of body composition use a two-component model of fat mass (FM) and fat-free mass (FFM). FFM is composed of the total body water mass, bone mineral mass, and protein mass, which collectively form a four-component model of body compositions (i.e., fat, water, protein, and bone), although the four-component model is used mainly in laboratory settings (226,227). The two-component model consisting of FM and FFM is the most easily applied body composition model in practice (33,227) and is used in this consensus paper. Bone mineral content and density (BMD) are both important components of body composition, as they are markers of bone health, with BMD having clinical relevance for the assessment of osteoporosis and fracture risk (228). Skeletal fitness was considered for inclusion as a subcomponent of body composition, but it was ultimately eliminated because there is a lack of valid and feasible assessment methods that can be used outside of a clinical setting at the current time (229–231).

Exercise-induced changes. Regular participation in exercise can change both FM and FFM. Several systematic reviews and scientific statements have established that clinically meaningful reductions in both total and abdominal FM are associated with exercise (232–240), with intra-abdominal or visceral FM being extremely responsive to exercise (241). Resistance exercise is recommended as the first-line therapy when increasing skeletal muscle mass is the primary objective and is appropriate to use throughout the lifespan (113,124,234,237,242,243). Indeed, resistance exercise can increase skeletal muscle mass independent of sex and age (45,111,234,244,245). Resistance exercise may also induce total and regional fat loss, particularly when combined with endurance training or as part of a multicomponent exercise regimen (45,237,239,246,247).

Ability to perform exercise. Body composition can both enhance and detract from the ability to perform exercise. Skeletal muscle mass represents a functional component of FFM in the body that is positively associated with strength and force production and, thus, improved capacity for physical work and performance. Conversely, FM is generally considered to be a nonfunctional component of body composition with increasing amounts of FM mechanically and metabolically hindering performance (59,248). Higher levels of body fat are associated with lower levels of VO_{2max} (249,250), and obesity is associated with lower physical function and PA (251).

Health-related benefits. A large body of evidence has established that the accumulation of excess FM is associated with increased risks of mortality and a wide range of morbidities (236,240,241,252–259), including elevated risks of developing cardiometabolic diseases, mental health disorders, some cancers, impaired postural stability, slower gait speed, and reduced physical function. However, it is now well established that excess accumulation of abdominal fat, especially visceral fat, and ectopic fat deposition in and around the liver, pancreas, and muscle (i.e., myosteatosis) is the obesity phenotype that conveys the greatest health risk and is associated with an increased incidence of type 2 diabetes, heart failure, stroke, kidney stones, and cancer (27,241,257–267). A sustained weight loss of 3–5% of initial body weight in

individuals with excess body weight and adiposity results in clinical meaningful reductions in cardiovascular disease risk factors (13).

Reduced skeletal muscle mass, with or without excess adiposity, is associated with higher risks for all-cause mortality, cardiovascular disease, and cancer (133,174,268–273). Skeletal muscle carries essential roles in metabolism that are particularly important relative to cardiometabolic diseases through regulation of glucose uptake, insulin responses, and fatty acid metabolism (274–276). Age- and inactivity-related losses of skeletal muscle mass are linked premature death (277–279) and to a substantially higher risk of adverse health outcomes, including increased risk of falls and fractures, impaired ability to perform ADLs, cardiopulmonary disease, cognitive impairment, and mobility disorders (280,281). Further, it contributes to lowered quality of life, loss of independence, and the need for long-term care (280,281).

Assessments. Methods to assess weight status and body composition that are the most feasible across health and fitness practice settings with respect to cost, expertise needed for quality measures, and acceptable reliability and validity include anthropometry, skinfolds, and bioelectrical impedance analysis (BIA) (237). Anthropometric measures including height, weight, hip circumference, and waist circumference can be used to calculate indices such as BMI, waist-to-height ratio, and waist-to-hip ratio to classify obesity-related health risks (237,282). The combination of BMI and waist circumference or BMI and the waist-to-height ratio identifies adults at increased health risk better than either measure alone (27,237,282,283). Although used widely, there are limitations to the application of BMI for the estimation of body fatness, especially in women, older individuals, and some people living with chronic diseases, and more precise assessments of body composition are recommended whenever possible (237,282–284).

Skinfolds can be used to estimate body fatness given the known association between subcutaneous fat and total body adiposity (13,227). BIA devices have gained popularity given the reductions in cost, portability, and measurement features. Typically, BIA devices measure the impedance, resistance, and reactance to the flow of an extremely low-strength electrical current passed through one or more body regions based on electrode configuration. Depending on the BIA system, measured components include not only FM and FFM but also fluid spaces such as total body, intracellular, and extracellular water and can estimate regional and total body adiposity and skeletal muscle (285,286).

Field methods (e.g., skinfolds, circumferences, and girths) can also be used to estimate skeletal muscle mass (287–290). Field measures have inherent limitations with respect to validity and reliability that should be considered when using these techniques in practice.

Neuromotor Fitness

Neuromotor fitness (Table 2) encompasses a variety of motor skills including locomotor, object manipulation, and stabilizing skills. These skills engage multiple body segments and form the foundation of physical activities, exercise, and recreational sports (291,292). Variables that have been proposed to fall under the construct of neuromotor fitness (previously termed skill-related fitness) are balance, agility, gait, coordination, speed of movement, and reaction time (9,11–13,15–19,21,22,24,62); however, a lack of precision in definitions and assessment techniques presents challenges to include all of these as subcomponents of neuromotor fitness. Currently, there is sufficient scientific evidence supporting each of the four inclusion criteria to include balance, agility, and gait speed as distinct subcomponents of neuromotor fitness in this consensus paper, as presented below. Coordination, speed of movement, and reaction time are recognized as important aspects of motor function and control—and they are identifiable parts of balance, gait, and agility—but, at the current time, there is insufficient evidence in adults supporting them as distinct subcomponents meeting all of the four inclusion criteria (293–298). For that reason, coordination, speed of movement, and reaction time are not considered separately and are not currently included in the updated model of PF, but they can be considered in the future as research evolves.

Exercise-induced changes. Neuromotor fitness can be enhanced by exercise training. Balance improves by participation in balance exercises (299–301), resistance exercise (301–303), core skeletal muscle strengthening (44), aerobic exercise (301,304), and dance (305,306). Balance exercises improve static and dynamic balance in healthy younger and older adults, and there is evidence of a dose–response association with greater frequency and duration of balance exercise practice (307,308). Studies of older adults have evaluated agility exercises alone (309), or as part of multicomponent exercise programs (309–312), and these training methods have improved agility, balance, gait speed, strength, physical function (e.g., chair rise), and working memory (309–313). Gait speed is improved by exercise training involving ambulation, resistance exercises, and multicomponent exercise (314–319). Multicomponent exercise programs, such as tai chi and yoga, and those that include combinations of exercise modalities (e.g., neuromotor, resistance, aerobic, or stretching exercises) result in improvements in several neuromotor fitness attributes (i.e., gait speed, balance, and agility), in addition to beneficial changes in muscular strength, body composition, flexibility, physical function, quality of life, and the risk of falling in younger and older adults, including those experiencing frailty (119,320–335).

Ability to perform exercise. Balance is a crucial component of movement control and stability for

activities ranging from the performance of ADLs to participation in recreational activities and sport (40,336–339). Maintaining stability during changes to the base of support (e.g., dynamic balance) during movement or while walking, running, and climbing stairs is critical in preventing falls and injuries that can lead to loss of mobility (340–344). Agility allows for smooth and efficient movement while changing direction and assists in maintaining proper alignment and balance during movement. While typically associated with sport, agility may also apply to other relevant movement domains such as functional mobility (143,311,313,345). There is an inverse relationship between agility and the fear of falling (346), which, in turn, is associated with a decrease in self-reported and performance-based physical function (310,311,346–348) and lower total daily PA (349,350).

Health-related benefits. Neuromuscular fitness is inversely associated with risks of cardiovascular diseases, hypertension, cognitive function (339,351–356), and psychological health (357,358). Neuromotor fitness and the subcomponents of balance, agility, and gait speed are associated with better health and functional outcomes in middle-aged and older adults (299,301,359–365).

Studies of exercise interventions including neuromotor and multicomponent exercises can improve ability to engage in ADLs, reduce frailty and depression, increase strength, improve cognition and quality of life, and reduce cardiovascular and inflammatory biomarkers in older adults (174,346,366–369). Interventions of multicomponent exercise programs of neuromotor exercise combined with aerobic exercise have resulted in improvements in cardiometabolic disease risk factors such as blood pressure, lipoprotein profile, vascular function, and glucose levels (370–372). However, there is a paucity of studies evaluating the physiological effects of neuromotor exercise alone.

Neuromotor exercises have been shown to result in changes of brain function and structure, and these adaptations are thought to underlie the observed changes in cognition, and executive and motor function (373). For example, activation of the posterior parietal network of the brain occurs with neuromotor exercise and this is associated with better movement activation, visual–spatial processing, and enhanced attention function (373–376). Additionally, the changes in the structure function of the motor areas of the brain (e.g., the cortico-striatal and the cortico-cerebellar systems) have been observed (373,376).

Impaired balance, agility, and gait are associated with elevated risks of falling (41,301,359,365,377,378). Declines in balance increase the risks of all-cause mortality, morbidity, and disability in middle-aged and older adults and in individuals with chronic diseases (362,365,377–380). Although many hypothesize that better agility may reduce musculoskeletal injuries, there is insufficient evidence in nonathletes and athletes to support this benefit (359,381). Agility training improves gait speed, lower limb power, static balance, and cognitive performance in older adults (311). A systematic review and meta-analysis of multimodal agility training revealed large overall effects on strength, mobility, and endurance and moderate effects on balance (312). Gait speed is linked to better health-related outcomes in adults ranging from 35 yr to older adults and in people living with disability. Slow gait speed is associated with increased risks of frailty, obesity, cognitive impairment, and increased cardiovascular disease morbidity and all-cause mortality (314,316,362,377,67,382–385).

Assessments. There is no one test to measure neuromotor fitness, but rather, a variety of tests are available to measure subcomponents of neuromotor fitness. Gait speed is measured as the time elapsed to walk a distance, such as in the 10-meter walk and 6-minute walk tests. Balance is measured in response to internal and external perturbations by static and dynamic tests such as the Functional Reach Test, Berg Balance Scale, Fullerton Advanced Balance Scale, Y Balance Test, and Four Stage Balance Test (386–391). Agility tests often measure only some aspects of agility, such as the physical component of change of direction and speed, or the cognitive components of anticipation and pattern recognition, with fewer comprehensive tests that include a contextual component (i.e., stimulus) (65,392). Measures of agility involving a change of direction or speed in response to a stimulus include the Timed Up and Go Test, various obstacle course tests, for example, Agility T-test and Illinois Agility Test (65,392). There is a need to expand agility testing to include more tests designed specifically for application to adults, because many neuromotor fitness tests are designed for athletes (65,392).

Flexibility

Flexibility (Table 2) refers to the range of motion (ROM) around a joint. It is a function of several factors that affect the skeletal muscles, joints, and tissues involved in moving the joint such as distensibility of the joint capsule, and tightness of ligaments and tendons, temperature, and viscoelastic properties of the skeletal muscle (393–395).

Exercise-induced changes. It is universally accepted that chronic stretch training will increase joint ROM (321,69,396). The beneficial effects of stretch training on flexibility are enhanced with higher frequency (number of days of exercise) and total duration of the stretching exercises (i.e., ≥5 min) (56). Recently, other effective strategies to increase flexibility, such as foam rolling (396,397) or resistance training (398), have been supported in the literature.

Ability to perform exercise. Higher levels of flexibility may contribute to improved performance of recreational sports including marathon running in women

(399) and golf (400). Adequate levels of flexibility may be important for the safe and effective participation of ADLs (401,402). For example, healthy middle-aged and older adults who participated in flexibility exercises were 24% less likely to report functional limitations compared with individuals who reported no stretch training (403). Flexibility is also associated with the ability to sit and rise from the floor in adults of all ages (404). However, flexibility typically declines with aging, and can be adversely affected by injuries, and a variety of diseases and conditions such as obesity, osteoarthritis, back pain, and inflammatory conditions (405–410). Impairments of joint ROM can lead to injuries, pain, and disability (403,405,411), which can compromise engagement in exercise (401,404,412–414). Participation in exercises that improve joint flexibility can reduce pain (401,405,415) improve physical functioning and facilitate engagement in PA (416–418).

Health-related benefits. Enhanced flexibility improves balance (401), and the risks of falling are decreased with better flexibility of the hip (419). A recent meta-analysis showed that stretching exercise interventions reduced the odds of muscular injuries by 37% without any significant differences in the odds of experiencing a tendinous injury in healthy adults (420). Stretch training reduces musculotendinous injuries in military personnel (421), or when engaging in activities involving explosive muscular contractions and changes of direction (401,402). Poor flexibility of the hamstring muscles has been proposed as a contributor to low back pain, but there is a dearth of high-quality research to determine if such a singular relationship between hamstrings flexibility and low back pain exists (422). However, two meta-analytical studies have supported that decreased hamstring flexibility and limited lateral flexion, along with lordosis of the lumbar spine, are associated with the development of low back pain (422,423), and Hafeti et al. showed that a program of stretching exercise reduced low back pain and improved ROM (405). Participation in stretch training improves arterial remodeling and vascular endothelial function and can reduce arterial stiffness, heart rate, and diastolic blood pressure (424–427). Stretch training has also been associated with favorable acute effects on several cardiovascular disease-related biomarkers, such as improved cardiac autonomic function and mucosal immune function (428–433). Although there is a paucity of studies examining the link between flexibility and all-cause mortality, a recent prospective study found an inverse relationship between flexibility and all-cause mortality in middle-aged men and women (434).

Assessments. There is a diverse array of instruments to directly and indirectly measure flexibility, including goniometers and digital levels (435–437), and functional tests such as the Schober and modified Schober Tests for measuring lumbar flexibility (438), the back scratch test (224), sit and reach (439–441), straight leg raise (442), and weight-bearing lunge test (443). Goniometers and digital levels have been evaluated as having high intrarater reliability (444,445), but they may not have consistently strong inter-rater (445) or between instrument reliability. Other assessments may not always measure what people expect. For example, the sit and reach test is commonly used to measure lower back and hamstrings flexibility (441,70,446–448); however, studies show mixed validity for different aspects of flexibility being measured, and factors like scapular abduction, arm and spine length differences, compensation strategies, and individual differences in discomfort perception can influence the results (449–451).

APPLICATION OF THE UPDATED MODEL OF PHYSICAL FITNESS

The expert panel proposed an updated model of PF that differs from the original model of the past decades in several ways. First, there was an effort to compile and present the components of PF in a clear and organized manner. The updated model of PF includes CRF, muscular fitness, body composition, neuromotor fitness, and flexibility as the major components, with muscular fitness, body composition, and neuromotor fitness further divided into subcomponents (Fig. 1). Muscular strength, muscular endurance, and muscular power were integrated into one PF component entitled muscular fitness, standardizing the categorization of this component and the related subcomponents. Second, the components of skill-related fitness were gathered into one category in a manner that is consistent with the other PF components and renamed neuromotor fitness. Third, the dichotomization of health-related and skill-related fitness has been eliminated because all components provide health and fitness benefits that are applicable to all adults. Moreover, consistency has been built into the updated model of PF as all components and subcomponents were clearly defined and met rigorous inclusion criteria (i.e., health benefits, can be changed by training, related to the ability to perform PA, measurable in practice), ensuring that there was sufficient scientific evidence to support their recognition as distinct aspects of PF. Finally, PF is conceptualized in a highly innovative way that emphasizes both interconnectedness and individualization as core tenets of the updated model PF as described below.

The interconnectedness among the components of PF was a consistent theme that emerged throughout this project (Fig. 2). The updated model moves beyond the traditional approach where the components are viewed in isolation to a model where PF is shown to be more complex with overlapping and interrelated attributes that may be responsive to a variety of exercise modalities.



FIGURE 1—Updated model of physical fitness.

This is an important revision that deserves emphasis and consideration in practice. While it is beyond the scope of this paper to outline all interrelationships, presented are some examples that emphasize the interconnectedness of fitness components. FFM, and its component of skeletal muscle mass, was included as a dimension of body composition. However, skeletal muscle hypertrophy is induced by resistance exercise and is one of the mechanisms whereby muscular fitness is improved. Additionally, there is evidence for small magnitude improvements in skeletal muscle hypertrophy with extensive stretch training to improve flexibility (416,452). Resistance exercise can result in improvements in both CRF and flexibility (398,453). Aerobic exercise training can improve balance (304,454) and gait speed (455,456) and attenuates age-related reductions in skeletal muscle strength (457). Multicomponent exercise programs (e.g., tai chi, Pilates, yoga, and dance), which include a combination of exercise modalities within a single program, improve balance and physical function, muscular strength, body composition, flexibility, agility, reaction time, and gait speed in younger and older adults, including those experiencing frailty (160,320,323,326–328,330–335,458). However, the common practice of combining exercises to improve fitness can make it difficult to identify the independent effects of the exercises involved, as the benefits can be synergistic or additive.

The concept of individualization emerged as an important consideration in this consensus statement. Rather than considering the components of PF and their related subcomponents as a static list of variables that apply to everyone in the same manner and to the same degree, they should be considered dynamic in that each can be highlighted or de-emphasized in the exercise prescription depending on the characteristics (e.g., health status, level of fitness, and disability) and priorities of the individual. For an older adult, it may be more important to participate in exercise to improve muscular fitness, neuromotor fitness, and body composition (FFM) because reducing risks of falling is a priority. Conversely, a younger adult might prioritize exercises to improve CRF and lower extremity strength to prepare, for example, for a community 5K road race. Further, an individual may have variable levels of fitness across the components, such that



FIGURE 2——The interconnectedness among the components of physical fitness.

they may have high fitness in one component and poor fitness in another, indicating the need for specific focus in weaker areas. Additionally, owing to genetic, health conditions, and other individual factors, not everyone is able to achieve the highest normative levels of fitness. While optimizing all components of fitness is ideal for a given individual, in practice, behavioral and psychological factors need to be considered for effective exercise programs to improve PF (459–461).

CONCLUSIONS

This consensus statement presents an updated model of PF, reflecting advances in science and recognizing the complexity of PF. It can be applied in practice directly to the individual as a tool to organize exercise program design and can continue to serve as a blueprint for PA guidelines (462). The updated model of PF is supported by scientific evidence and each component of PF included in the model meets four criteria (i.e., can be changed by exercise, impacts the ability to perform exercise and PA; contributes to health; can be feasibly assessed in practice). Importantly, the updated model standardizes the definitions of PF and its components, and it leaves room to add new components and subcomponents over time as additional evidence or assessment technologies come to light.

The authors thank Jimin Kim who provided the graphic designs included in this paper. The results of this study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of the present study do not constitute endorsement by the American College of Sports Medicine.

No funding or conflicts of interest were disclosed.

REFERENCES

1. Appelqvist-Schmidlechner K, Vaara JP, Vasankari T, Häkkinen A, Mäntysaari M, Kyröläinen H. Muscular and cardiorespiratory fitness are associated with health-related quality of life among young adult men. *BMC Public Health*. 2020;20(1):842.
2. Cremer PC, Wu Y, Ahmed HM, et al. Use of sex-specific clinical and exercise risk scores to identify patients at increased risk for all-cause mortality. *JAMA Cardiol*. 2017;2(1):15–22.
3. Evaristo S, Moreira C, Lopes L, et al. Muscular fitness and cardiorespiratory fitness are associated with health-related quality of life: Results from labmed PA study. *J Exerc Sci Fit*. 2019;17(2):55–61.
4. Lang JJ, Prince SA, Merucci K, et al. Cardiorespiratory fitness is a strong and consistent predictor of morbidity and mortality among adults: An overview of meta-analyses

representing over 20.9 million observations from 199 unique cohort studies. *Br J Sports Med*. 2024;58(10):556–66.
5. Lee J. The effects of resistance training on muscular strength and hypertrophy in elderly cancer patients: A systematic review and meta-analysis. *J Sport Health Sci*. 2022;11(2):194–201.
6. Mandsager K, Harb S, Cremer P, Phelan D, Nissen SE, Jaber W. Association of cardiorespiratory fitness with long-term mortality among adults undergoing exercise treadmill testing. *JAMA Netw Open*. 2018;1(6):e183605.
7. Pedersen BK, Saltin B. Exercise as medicine—evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports*. 2015;25(Suppl 3):1–72.
8. Ross R, Blair SN, Arena R, et al; American Heart Association Physical Activity Committee of the Council on Lifestyle and Cardiometabolic Health. Importance of assessing cardiorespiratory fitness in clinical practice: A case for fitness as a clinical vital sign: A scientific statement from the American Heart Association. *Circulation*. 2016;134(24):e653–99.
9. U. S. Department of Health and Human Services. *2018 Physical Activity Guidelines Advisory Committee Scientific Report*. U.S. Department of Health and Human Services; 2018. Available from: https://health.gov/sites/default/files/2019-09/PAG_Advisory_Committee_Report.pdf.
10. Warburton DE, Glendhill N, Quinney A. The effects of changes in musculoskeletal fitness on health. *Can J Appl Physiol*. 2001;26(2):161–216.
11. Garber CE, Blissmer B, Deschenes MR, et al; American College of Sports Medicine. American College of Sports Medicine Position Stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;43(7):1334–59.
12. Liguori G, editor. *ACSM's Guidelines for Exercise Testing and Prescription 11th Edition*. Eleventh edition. Wolters Kluwer; 2021.
13. Ozemek C, editor. *ACSM's Guidelines for Exercise Testing and Prescription 12th Edition*. 12th ed. Wolters Kluwer; 2025.
14. Pate RR. A new definition of youth fitness. *Phys Sportsmed*. 1983;11(4):77–83.
15. U.S. Department of Health and Human Services. *2008 Physical Activity Guidelines Advisory Committee Scientific Report*. Department of Health and Human Services; 2008. Available from: https://odphp.health.gov/sites/default/files/2019-10/CommitteeReport_7.pdf.
16. Franklin BA, editor. *ACSM's Guidelines for Exercise Testing and Prescription 6th Edition*. 6th ed. Lippincott Williams & Wilkins; 2000.
17. Kenney WL, editor. *ACSM's Guidelines for Exercise Testing and Prescription 5th Edition*. 5th ed. Williams & Wilkins; 1995.
18. Pescatello LS, editor. *ACSM's Guidelines for Exercise Testing and Prescription 9th Edition*. 9th ed. Wolters Kluwer/Lippincott Williams & Wilkins Health; 2014.
19. Riebe D, editor. *ACSM's Guidelines for Exercise Testing and Prescription 10th Edition*. 10th ed. Wolters Kluwer/Lippincott Williams & Wilkins Health; 2018.
20. Thompson PD, editor. *ACSM's Guidelines for Exercise Testing and Prescription 3rd Edition*. 3rd ed. Lea & Febiger; 1986.
21. Thompson WR, editor. *ACSM's Guidelines for Exercise Testing and Prescription 8th Edition*. 8th ed. Wolters Kluwer/Lippincott Williams & Wilkins Health; 2010.
22. Whaley MH, editor. *ACSM's Guidelines for Exercise Testing and Prescription 7th Edition*. 7th ed. Lippincott Williams & Wilkins; 2006. p. xxi, 366 ill. 22 cm.
23. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Rep*. 1985;100(2):126–31.
24. U. S. Department of Health and Human Services. *Physical Activity and Health: A Report of the Surgeon General*. Centers for Disease Control and Prevention (US), National Center for Chronic Disease Prevention and Health Promotion; 1996. .
25. Adams M, Gordt-Oesterwind K, Bongartz M, et al. Effects of physical activity interventions on strength, balance and falls in middle-aged adults: A systematic review and meta-analysis. *Sports Med Open*. 2023;9(1):61.
26. Australian Department of Health. Physical activity and exercise. *Physical Activity and Exercise*. 2026; [cited Jan 8, 2026]. Available from: https://www.health.gov.au/topics/physical-activity-and-exercise.
27. Ross R, Chaput J, Giangregorio LM. Canadian 24-hour movement guidelines for adults aged 18–64 years and adults aged 65 years or older: An integration of physical activity, sedentary behaviour, and sleep. *Appl Physiol Nutr Metab [Internet]*. 2020;45(10 Suppl. 2):S57–S102.
28. World Health Organization. *WHO Guidelines on Physical Activity and Sedentary Behaviour*. World Health Organization; 2020.
29. American College of Sports Medicine. Roundtables & Specialty Conferences. 2026; [cited 2026 Jan 9] Available from: https://acsm.org/meetings/specialty/.
30. Blazey P, Crossley KM, Ardern CL, Middelkoop M, Scott A, Khan KM. It is time for consensus on "consensus statements." *Br J Sports Med*. 2022;56(6):306–7.
31. Aparicio Rodrigo M, González Rodríguez P, Balado Insunza N, Fraile Astorga G, Aizpurua Galdeano P, Ochoa Sangrador C; Comité de Trabajo de Pediatría Basada en la Evidencia. How to develop and evaluate consensus documents: Methods and checklists. *An Pediatr (Engl Ed)*. 2025;103(1):503890.
32. Rosenfeld RM, Nnacheta LC, Corrigan MD. Clinical consensus statement development manual. *Otolaryngol Head Neck Surg*. 2015;153(2 Suppl):S1–S14.
33. Bishop DJ, Beck B, Biddle SJH, et al. Physical activity and exercise intensity terminology: A joint american college of sports medicine (ACSM) expert statement and exercise and sport science australia (ESSA) consensus statement. *J Sci Med Sport*. 2025;28(12):980–91.
34. Campbell N, De Jesus S, Prapavessis H. Physical Fitness. In: Gellman MD, Turner JR, editors. *Encyclopedia of Behavioral Medicine*. Springer; 2013. pp. 1486–9.
35. Corbin CB, Le Masurier G. *Fitness For Life*. 6th ed. Human Kinetics; 2014.
36. Edwards P, Hauser J. Fitness. *Canadian Encyclopedia*. 2015; [cited Jan 8, 2026]. Available from: https://thecanadianencyclopedia.ca/en/article/fitness.
37. Merriam-Webster. "Physical fitness. *Merriam-Webster.com Dictionary*. 2026; Available from: https://www.merriam-webster.com/dictionary/physical%20fitness.
38. Warburton DER, Nicol CW, Bredin SSD. Health benefits of physical activity: The evidence. *CMAJ*. 2006;174(6):801–9.
39. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451–62.
40. Alhwoaimel NA, Alshehri MM, Alhowimel AS, Alenazi AM, Alqahtani BA. Functional mobility and balance confidence measures are associated with disability among community-dwelling older adults. *Medicina*. 2024;60(9):1549.
41. Morat M, Faude O, Hanssen H, et al. Agility training to integratively promote neuromuscular, cognitive, cardiovascular and psychosocial function in healthy older adults: A study protocol of a one-year randomized-controlled trial. *Int J Environ Res Public Health [Internet]*. 2020;17(6):1853.
42. Orr R, de Vos NJ, Singh NA, Ross DA, Stavrinos TM, Fiatarone-Singh MA. Power training improves balance in healthy older adults. *J Gerontol A Biol Sci Med Sci*. 2006;61(1):78–85.

43. Shih Y-L, Shih C-C, Chen J-Y. The association between walking speed and risk of cardiovascular disease in middle-aged and elderly people in Taiwan, a community-based, cross-sectional study. *PLoS One*. 2020;15(7):e0235277.
44. Granacher U, Gollhofer A, Hortobágyi T, Kressig RW, Muehlbauer T. The importance of trunk muscle strength for balance, functional performance, and fall prevention in seniors: A systematic review. *Sports Med*. 2013;43(7):627–41.
45. Lopez P, Taaffe DR, Galvão DA, et al. Resistance training effectiveness on body composition and body weight outcomes in individuals with overweight and obesity across the lifespan: A systematic review and meta-analysis. *Obes Rev*. 2022;24:e13428.
46. Phillips SM, Ma JK, Rawson ES. The coming of age of resistance exercise as a primary form of exercise for health. *ACSM'S Health Fit J*. 2023;27(6):19–25.
47. Porcari JP, Bryant CX, Comana F, editors. *Exercise Physiology*. F. A. Davis Company; 2015.
48. Lee J, Zhang XL. Physiological determinants of VO2max and the methods to evaluate it: A critical review. *Sci Sports*. 2021;36(4):259–71.
49. Perry AS, Farber-Eger E, Gonzales T, et al. Proteomic analysis of cardiorespiratory fitness for prediction of mortality and multisystem disease risks. *Nat Med*. 2024;30(6):1711–21.
50. Anaby D, Miller WC, Eng JJ, Jarus T, Noreau L; PACC Research Group. Can personal and environmental factors explain participation of older adults? *Disabil Rehabil*. 2009;31(15):1275–82.
51. Arena R, Pronk NA, Woodard C. The influence of social vulnerability and culture on physical inactivity in the united states—identifying hot spots in need of attention. *Am J Med*. 2024;137(2):113–21.
52. Barnett DW, Barnett A, Nathan A, Van Cauwenberg J, Cerin E; Council on Environment and Physical Activity (CEPA)—Older Adults working group. Built environmental correlates of older adults' total physical activity and walking: A systematic review and meta-analysis. *Int J Behav Nutr Phys Act*. 2017;14(1):103.
53. Seefeldt V, Malina RM, Clark MA. Factors affecting levels of physical activity in adults. *Sports Med*. 2002;32(3):143–68.
54. Fraser BJ, Huynh QL, Schmidt MD, Dwyer T, Venn AJ, Magnussen CG. Childhood muscular fitness phenotypes and adult metabolic syndrome. *Med Sci Sports Exerc*. 2016;48(9):1715–22.
55. Smith JJ, Eather N, Morgan PJ, Plotnikoff RC, Faigenbaum AD, Lubans DR. The health benefits of muscular fitness for children and adolescents: a systematic review and meta-analysis. *Sports Med*. 2014;44(9):1209–23.
56. Thomas E, Bianco A, Raia T, et al. Relationship between velocity and muscular endurance of the upper body. *Hum Mov Sci*. 2018;60:175–82.
57. Johnson D, Lynch J, Nash K, Cygan J, Mayhew JL. Relationship of lat-pull repetitions and pull-ups to maximal lat-pull and pull-up strength in men and women. *J Strength Cond Res*. 2009;23(3):1022–8.
58. Cormie P, McGuigan MR, Newton RU. Developing maximal neuromuscular power: Part 2—training considerations for improving maximal power production. *Sports Med*. 2011;41(2):125–46.
59. Ackland TR, Lohman TG, Sundgot-Borgen J, et al. Current status of body composition assessment in sport: Review and position statement on behalf of the ad hoc research working group on body composition health and performance, under the auspices of the I.O.C. Medical Commission. *Sports Med*. 2012;42(3):227–49.
60. Heymsfield SB, Brown J, Ramirez S, Prado CM, Tinsley GM, Gonzalez MC. Are lean body mass and fat-free mass the same or different body components? A critical perspective. *Adv Nutr*. 2024;15(12):100335.
61. Holmes CJ, Racette SB. The utility of body composition assessment in nutrition and clinical practice: An overview of current methodology. *Nutrients*. 2021;13(8):2493.
62. Stodden DF, Goodway JD, Langendorfer SJ, et al. A developmental perspective on the role of motor skill competence in physical activity: An emergent relationship. *Quest*. 2008;60(2):290–306.
63. Barnett LM, Lai SK, Veldman SLC, et al. Correlates of gross motor competence in children and adolescents: A systematic review and meta-analysis. *Sports Med*. 2016;46(11):1663–88.
64. King-Dowling S, Proudfoot NA, Cairney J, Timmons BW. Motor competence, physical activity, and fitness across early childhood. *Med Sci Sports Exerc*. 2020;52(11):2342–8.
65. Sheppard JM, Young WB. Agility literature review: Classifications, training and testing. *J Sports Sci*. 2006;24(9):919–32.
66. Pollock AS, Durward BR, Rowe PJ, Paul JP. What is balance? *Clin Rehabil*. 2000;14(4):402–6.
67. Mehmet H, Robinson SR, Yang AWH. Assessment of gait speed in older adults. *J Geriatr Phys Ther*. 2020;43(1):42–52.
68. Middleton A, Fritz SL, Lusardi M. Walking speed: The functional vital sign. *J Aging Phys Act*. 2015;23(2):314–22.
69. Behm DG. *The Science and Physiology of Flexibility and Stretching: Implications and Applications in Sport Performance and Health*. Routledge Publishers; 2018.
70. Liemohn W. Flexibility and muscular strength. *J Phys Educ Recreat Dance*. 1988;59(7):37–40.
71. McHugh MP, Kremenic IJ, Fox MB, Gleim GW. The role of mechanical and neural restraints to joint range of motion during passive stretch. *Med Sci Sports Exerc*. 1998;30(6):928–32.
72. Ross R, Myers J. Cardiorespiratory fitness and its place in medicine. *Rev Cardiovasc Med*. 2023;24(1):14.
73. Poole DC, Jones AM. Measurement of the maximum oxygen uptake Vo2max: Vo2peak is no longer acceptable. *J Appl Physiol (1985)*. 2017;122(4):997–1002.
74. Chase JA, Conn V. Meta-analysis of fitness outcomes from motivational physical activity interventions. *Nurs Res*. 2013;62(5):294–304.
75. Edwards JJ, Deenmamode AHP, Griffiths M, et al. Exercise training and resting blood pressure: A large-scale pairwise and network meta-analysis of randomised controlled trials. *Br J Sports Med*. 2023;57(20):1317–26.
76. Edwards JJ, Griffiths M, Deenmamode AHP, O'Driscoll JM. High-intensity interval training and cardiometabolic health in the general population: A systematic review and meta-analysis of randomised controlled trials. *Sports Med*. 2023;53(9):1753–63.
77. Maurits D, Achmad Bayu RF, Hsing Mei C. Physical activity improves health-related quality of life, 6MWT, and VO(2) peak before and during COVID-19 in patients with heart failure: A meta-analysis. *Semergen*. 2023;49(8):102039.
78. Poon ET, Li HY, Gibala MJ, Wong SH, Ho RS. High-intensity interval training and cardiorespiratory fitness in adults: An umbrella review of systematic reviews and meta-analyses. *Scand J Med Sci Sports*. 2024;34(5):e14652.
79. Franklin BA, Eijsvogels TMH, Pandey A, Quindry J, Toth PP. Physical activity, cardiorespiratory fitness, and cardiovascular health: A clinical practice statement of the American society for preventive cardiology part II: Physical activity, cardiorespiratory fitness, minimum and goal intensities for exercise training, prescriptive methods, and special patient populations. *Am J Prev Cardiol*. 2022;12:100425.
80. Zeiher J, Ombrellaro KJ, Perumal N, Keil T, Mensink GBM, Finger JD. Correlates and determinants of cardiorespiratory fitness in adults: A systematic review. *Sports Med Open*. 2019;5(1):39.
81. Hellsten Y, Nyberg M. Cardiovascular adaptations to exercise training. *Compr Physiol*. 2015;156(1):1–32.
82. Herrmann SD, Willis EA, Ainsworth BE, et al. Adult compendium of physical activities: A third update of the energy costs of human activities. *J Sport Health Sci*. 2024;13(1):6–12.

83. Willis EA, Herrmann SD, Hastert M, et al. Older adult compendium of physical activities: Energy costs of human activities in adults aged 60 and older. *J Sport Health Sci*. 2024;13(1):13–7.
84. Warner A, Vanicek N, Benson A, Myers T, Abt G. Agreement and relationship between measures of absolute and relative intensity during walking: A systematic review with meta-regression. *PLoS One*. 2022;17(11):e0277031.
85. Yu R, Lai D, Leung G, Tong C, Woo J. Using cardiorespiratory fitness to operationalize vitality: A path analysis on the hierarchical structure of intrinsic capacity. *J Nutr Health Aging*. 2024;28(8):100300.
86. Vanzella LM, Ribeiro F, Laurino MJL, et al. Unveiling the link between physical parameters and safety in cardiac rehabilitation: Longitudinal observational study: Physical parameters and cardiac adverse events. *Curr Probl Cardiol*. 2025;50(1):102916.
87. Singh B, Cadenas-Sanchez C, Costa BGG, et al. Comparison of objectively measured and estimated cardiorespiratory fitness to predict all-cause and cardiovascular disease mortality in adults: A systematic review and meta-analysis of 42 studies representing 35 cohorts and 3.8 million observations. *J Sport Health Sci*. 2025;14:100986.
88. Franklin BA, Wedig IJ, Sallis RE, Lavie CJ, Elmer SJ. Physical activity and cardiorespiratory fitness as modulators of health outcomes: A compelling research-based case presented to the medical community. *Mayo Clin Proc*. 2023;98(2):316–31.
89. Kokkinos P, Faselis C, Samuel IBH, et al. Cardiorespiratory fitness and mortality risk across the spectra of age, race, and sex. *J Am Coll Cardiol*. 2022;9(6):80598–609.
90. Harber MP, Kaminsky LA, Arena R, et al. Impact of cardiorespiratory fitness on all-cause and disease-specific mortality: Advances since 2009. *Prog Cardiovasc Dis*. 2017;60(1):11–20.
91. Al-Mallah MH, Sakr S, Al-Qunaibet A. Cardiorespiratory fitness and cardiovascular disease prevention: An update. *Curr Atheroscler Rep*. 2018;20(1):1.
92. de Figueiredo JA, Garcia MC, Oliveira AJ. Relationship between cardiorespiratory fitness and falls in older adults: A population-based study. *Geriatr Gerontol Int*. 2025;25(2):167–72.
93. Myers J, Cadenas-Sanchez C, Ross R, Kokkinos P. The critical role of cardiorespiratory fitness in disease prevention. *J Sports Med Phys Fitness*. 2024;64(12):1361–71.
94. Ross R, Arena R, Myers J, Kokkinos P, Kaminsky LA. Update to the 2016 american heart association cardiorespiratory fitness statement. *Prog Cardiovasc Dis*. 2024;83:10–5.
95. Alfitian J, Riedel B, Ismail H, et al; METS Study Investigators. Sex-related differences in functional capacity and its implications in risk stratification before major non-cardiac surgery: A post hoc analysis of the international METS study. *EClinicalMedicine*. 2023;64:102223.
96. de Souza E Silva CG, Buginga GC, de Souza E Silva EA, et al. Prediction of mortality in coronary artery disease: Role of machine learning and maximal exercise capacity. *Mayo Clin Proc*. 2022;97(8):1472–82.
97. Tan S, Thang YW, Mulley WR, et al. Prognostic value of exercise capacity in kidney transplant candidates. *J Am Heart Assoc*. 2022;11(12):e025862.
98. Wang S, Xu L, Yang W, et al. Association of cardiorespiratory fitness with dementia risk across different levels of genetic predisposition: A large community-based longitudinal study. *Br J Sports Med*. 2025;59(3):150–8.
99. Kandola A, Ashdown-Franks G, Stubbs B, Osborn DPJ, Hayes JF. The association between cardiorespiratory fitness and the incidence of common mental health disorders: A systematic review and meta-analysis. *J Affect Disord*. 2019;257:748–57.
100. Schuch FB, Vancampfort D, Sui X, et al. Are lower levels of cardiorespiratory fitness associated with incident depression? A systematic review of prospective cohort studies. *Prev Med*. 2016;93:159–65.
101. Dores H, Mendes M, Abreu A, et al. Cardiopulmonary exercise testing in clinical practice: Principles, applications, and basic interpretation. *Rev Port Cardiol*. 2024;43(9):525–36.
102. Chaudhry S, Kumar N, Arena R, Verma S. The evolving role of cardiopulmonary exercise testing in ischemic heart disease—state of the art review. *Curr Opin Cardiol*. 2023;38(6):552–72.
103. Bonikowske AR, Taylor JL, Larson KF, et al. Evaluating current assessment techniques of cardiorespiratory fitness. *Expert Rev Cardiovasc Ther*. 2024;22(6):231–41.
104. Balady GJ, Williams MA, Ades PA, et al; American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: A scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. 2007;115(20):2675–82.
105. Fletcher GF, Ades PA, Kligfield P, et al; American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee of the Council on Clinical Cardiology, Council on Nutrition, Physical Activity and Metabolism, Council on Cardiovascular and Stroke Nursing, and Council on Epidemiology and Prevention. Exercise standards for testing and training: A scientific statement from the American Heart Association. *Circulation*. 2013;128(8):873–934.
106. Sartor F, Vernillo G, Morree HM, et al. Estimation of maximal oxygen uptake via submaximal exercise testing in sports, clinical, and home settings. *Sports Med*. 2013;43(9):865–73.
107. Henriques J, Carvalho P, Rocha T, et al. A non-exercise based V02max prediction using FRIEND dataset with a neural network. *Annu Int Conf IEEE Eng Med Biol Soc*. 2017;2017:4203–6.
108. Peterman JE, Whaley MH, Harber MP, et al. Comparison of non-exercise cardiorespiratory fitness prediction equations in apparently healthy adults. *Eur J Prev Cardiol*. 2021;28(2):142–148–48.
109. Schumacher BT, Di C, Bellettiere J, et al. Validation, recalibration, and predictive accuracy of published VO2max prediction equations for adults ages 50-96 Yr. *Med Sci Sports Exerc*. 2023;55(2):322–32.
110. Björkman F, Ekblom O, Ekblom-Bak E, Bohman T. The ability of a submaximal cycle ergometer test to detect longitudinal changes in VO2max. *BMC Sports Sci Med Rehabil*. 2021;13(1):156.
111. American College of Sports Medicine. American college of sports medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc*. 2009;41(3):687–708.
112. Ferná Ortega JA, Mendoza Romero D, Sarmento H, Prieto Mondragón L, Rodríguez Buitrago JA. Relationship between dynamic and isometric strength, power, speed, and average propulsive speed of recreational athletes. *J Funct Morphol Kinesiol*. 2022;7(4):79.
113. Fragala MS, Cadore EL, Dorgo S, et al. Resistance training for older adults: Position statement from the national strength and conditioning association. *J Strength Cond Res*. 2019;33(8):2019–52.

114. Naclerio FJ, Colado JC, Rhea MR, Bunker D, Triplett NT. The influence of strength and power on muscle endurance test performance. *J Strength Cond Res*. 2009;23(5):1482–8.
115. Hughes DC, Ellefsen S, Baar K. Adaptations to endurance and strength training. *Cold Spring Harb Perspect Med*. 2018;8(6):a029769.
116. Knuttgen HG, Kraemer WJ. Terminology and measurement in exercise performance. *J Strength Cond Res*. 1987;1(1):1–10.
117. Knowles KS, Pagan JI, Beausejour JP, et al. Changes in muscle quality following short-term resistance training in older adults: A comparison of echo intensity and texture analysis. *Ultrasound Med Biol*. 2025;51(4):675–82.
118. Nunes JP, Blazevich AJ, Schoenfeld BJ, et al. Determining changes in muscle size and architecture after exercise training: One site does not fit all. *J Strength Cond Res*. 2024;38(4):787–90.
119. Vieira FT, Cai Y, Gonzalez MC, Goodpaster BH, Prado CM, Haqq AM. Poor muscle quality: A hidden and detrimental health condition in obesity. *Rev Endocr Metab Disord*. 2025;26(5):723–44.
120. Ghasemi Kahrizsangi N, Faridi Dastjerdi MA, Golafshan F, Ghasemi Kahrizsangi N, Shafieyoun MH, Parsaeefar MH. Resistance training and blood pressure: Insights from a meta-analysis of the aging population. *Arch Gerontol Geriatr*. 2025;137:105941.
121. McIntosh MC, Anglin DA, Robinson AT, Beck DT, Roberts MD. Making the case for resistance training in improving vascular function and skeletal muscle capillarization. *Front Physiol*. 2024;15:1338507.
122. Bagheri N, Bagheri R, Mesinovic J, et al. Effects of resistance training on muscular adaptations and inflammatory markers in overweight and obese men. *Med Sci Sports Exerc*. 2025;57(3):600–12.
123. Murach KA, Bagley JR. A primer on global molecular responses to exercise in skeletal muscle: Omics in focus. *J Sport Health Sci*. 2025;15:101029.
124. Carvalho L, Junior RM, Barreira J, Schoenfeld BJ, Orazem J, Barroso R. Muscle hypertrophy and strength gains after resistance training with different volume-matched loads: A systematic review and meta-analysis. *Appl Physiol Nutr Metab*. 2022;47(4):357–68.
125. Currier BS, Mcleod JC, Banfield L, et al. Resistance training prescription for muscle strength and hypertrophy in healthy adults: A systematic review and Bayesian network meta-analysis. *Br J Sports Med*. 2023;57(18):1211–20.
126. El Hadouchi M, Kiers H, de Vries R, Veenhof C, van Dieën J. Effectiveness of power training compared to strength training in older adults: A systematic review and meta-analysis. *Eur Rev Aging Phys Act*. 2022;19(1):18.
127. Mcleod JC, Currier BS, Lowisz CV, Phillips SM. The influence of resistance exercise training prescription variables on skeletal muscle mass, strength, and physical function in healthy adults: An umbrella review. *J Sport Health Sci*. 2024;13(1):47–60.
128. Mikkonen RS, Ihalainen JK, Hackney AC, Häkkinen K. Perspectives on concurrent strength and endurance training in healthy adult females: A systematic review. *Sports Med (Auckland)* 2024;54(3):673–96.
129. Peterson MD, Rhea MR, Alvar BA. Applications of the dose-response for muscular strength development: A review of meta-analytic efficacy and reliability for designing training prescription. *J Strength Cond Res*. 2005;19(4):950–8.
130. Rossi C, Vasiljevic I, Manojlovic M, et al. Optimizing strength training protocols in young females: A comparative study of velocity-based and percentage-based training programs. *Heliyon*. 2024;10(9):e30644.
131. Sarabia JM, Moya-Ramón M, Hernández-Davó JL, Fernandez-Fernandez J, Sabido R. The effects of training with loads that maximise power output and individualised repetitions vs. traditional power training. *PLoS One*. 2017;12(10):e0186601.
132. Schoenfeld B, Fisher J, Grgic J, et al. Resistance training recommendations to maximize muscle hypertrophy in an athletic population: Position stand of the IUSCA. *Int'l J Strength Cond*. 2021;1(1):1–30. [cited Sept 26, 2021]. Available from: https://journal.iusca.org/index.php/Journal/article/view/81
133. Wang Y, Luo D, Liu J, Song Y, Jiang B, Jiang H. Low skeletal muscle mass index and all-cause mortality risk in adults: A systematic review and meta-analysis of prospective cohort studies. *PLoS One*. 2023;18(6):e0286745.
134. Peterson MD, Pistilli E, Haff GG, Hoffman EP, Gordon PM. Progression of volume load and muscular adaptation during resistance exercise. *Eur J Appl Physiol*. 2011;111(6):1063–71.
135. Schoenfeld BJ, Peterson MD, Ogborn D, Contreras B, Sonmez GT. Effects of low- vs. high-load resistance training on muscle strength and hypertrophy in well-trained men. *J Strength Cond Res*. 2015;29(10):2954–63.
136. Foldvari M, Clark M, Laviolette LC, et al. Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A Biol Sci Med Sci*. 2000;55(4):M192–199.
137. Miszko TA, Cress ME, Slade JM, Covey CJ, Agrawal SK, Doerr CE. Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci*. 2003;58(2):171–5.
138. Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic performance. *Sports Med*. 2016;46(10):1419–49.
139. Eihara Y, Takao K, Sugiyama T, et al. Heavy resistance training versus plyometric training for improving running economy and running time trial performance: A systematic review and meta-analysis. *Sports Med Open*. 2022;8(1):138.
140. Makaruk H, Starzak M, Tarkowski P, Sadowski J, Winchester J. The effects of resistance training on sport-specific performance of elite athletes: A systematic review with meta-analysis. *J Hum Kinet*. 2024;91(ue):135–55.
141. Torres-Ronda L, Sánchez-Medina L, González-Badillo JJ. Muscle strength and golf performance: A critical review. *J Sports Sci Med*. 2011;10(1):9–18.
142. Ehlert A. The effects of strength and conditioning interventions on golf performance: A systematic review. *J Sports Sci*. 2020;38(23):2720–31.
143. Myers B, Hanks J. Hip strength, change of direction, and falls in recreational pickleball players. *Int J Sports Phys Ther*. 2024;19(9):1116–25.
144. Thiele D, Prieske O, Chaabene H, Granacher U. Effects of strength training on physical fitness and sport-specific performance in recreational, sub-elite, and elite rowers: A systematic review with meta-analysis. *J Sports Sci*. 2020;38(10):1186–95.
145. Mikkola J, Vesterinen V, Taipale R, Capostagno B, Häkkinen K, Nummela A. Effect of resistance training regimens on treadmill running and neuromuscular performance in recreational endurance runners. *J Sports Sci*. 2011;29(13):1359–71.
146. Orange ST, Madden LA, Vince RV. Resistance training leads to large improvements in strength and moderate improvements in physical function in adults who are overweight or obese: A systematic review. *J Physiother*. 2020;66(4):214–24.
147. Keating CJ, Cabrera-Linares JC, Párraga-Montilla JA, Latorre-Román PA, Del Castillo RM, García-Pinillos F. Influence of resistance training on gait & balance parameters in older adults: A systematic review. *Int J Environ Res Public Health*. 2021;18(4):1759.

148. Bårdstu HB, Andersen V, Fimland MS, et al. Effectiveness of a resistance training program on physical function, muscle strength, and body composition in community-dwelling older adults receiving home care: A cluster-randomized controlled trial. *Eur Rev Aging Phys Act*. 2020;17:11.
149. Liu C, Latham N. Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev*. 2009;2009(3):CD002759. doi:10.1002/14651858.CD002759.pub2.
150. Borges-Silva F, Martínez-Pascual M, Colomer-Poveda D, Márquez G, Romero-Arenas S. Does heavy-resistance training improve mobility and perception of quality of life in older women? *Biology (Basel)*. 2022;11(5):626.
151. Santos L, Ribeiro AS, Schoenfeld BJ, et al. The improvement in walking speed induced by resistance training is associated with increased muscular strength but not skeletal muscle mass in older women. *Eur J Sport Sci*. 2017;17(4):488–94.
152. Olsen PO, Termannsen AD, Bramming M, Tully MA, Caserotti P. Effects of resistance training on self-reported disability in older adults with functional limitations or disability—a systematic review and meta-analysis. *Eur Rev Aging Phys Act [Internet]*. 2019;16(24):24.
153. Soysal P, Hurst C, Demurtas J, et al. Handgrip strength and health outcomes: Umbrella review of systematic reviews with meta-analyses of observational studies. *J Sport Health Sci*. 2021;10(3):290–5.
154. Shailendra P, Baldock KL, Li LSK, Bennie JA, Boyle T. Resistance training and mortality risk: A systematic review and meta-analysis. *Am J Prev Med*. 2022;63(2):277–85.
155. Saeidifard F, Medina-Inojosa JR, West CP, et al. The association of resistance training with mortality: A systematic review and meta-analysis. *Eur J Prev Cardiol*. 2019;26(15):1647–65.
156. Stamatakis E, Lee IM, Bennie J, et al. Does strength-promoting exercise confer unique health benefits? A pooled analysis of data on 11 population cohorts with all-cause, cancer, and cardiovascular mortality endpoints. *Am J Epidemiol*. 2018;187(5):1102–12.
157. Tarasenko YN, Linder DF, Miller EA. Muscle-strengthening and aerobic activities and mortality among 3+ year cancer survivors in the U.S. *Cancer Causes Control*. 2018;29(4-5):475–84.
158. Ashton RE, Tew GA, Aning JJ, Gilbert SE, Lewis L, Saxton JM. Effects of short-term, medium-term and long-term resistance exercise training on cardiometabolic health outcomes in adults: Systematic review with meta-analysis. *Br J Sports Med*. 2020;54(6):341–8.
159. Costa RR, Buttelli ACK, Vieira AF, et al. Effect of strength training on lipid and inflammatory outcomes: Systematic review with meta-analysis and meta-regression. *J Phys Act Health*. 2019;16(6):477–91.
160. Huang L, Fang Y, Tang L. Comparisons of different exercise interventions on glycemic control and insulin resistance in prediabetes: A network meta-analysis. *BMC Endocr Disord*. 2021;21(1):181.
161. Lee J, Kim D, Kim C. Resistance Training for glycemic control, muscular strength, and lean body mass in old type 2 diabetic patients: A meta-analysis. *Diabetes Ther*. 2017;8(3):459–73.
162. Lemes IR, Ferreira PH, Linares SN, Machado AF, Pastre CM, Netto J. Resistance training reduces systolic blood pressure in metabolic syndrome: A systematic review and meta-analysis of randomized controlled trials. *Br J Sports Med*. 2016;50(23):1438–42.
163. MacDonald HV, Johnson BT, Huedo-Medina TB, et al. Dynamic resistance training as stand-alone antihypertensive lifestyle therapy: A meta-analysis. *J Am Heart Assoc*. 2016;5(10):e003231.
164. Correia RR, Veras ASC, Tebar WR, Rufino JC, Batista VRG, Teixeira GR. Strength training for arterial hypertension treatment: A systematic review and meta-analysis of randomized clinical trials. *Sci Rep*. 2023;13(1):201.
165. Gordon BR, McDowell CP, Lyons M, Herring MP. The effects of resistance exercise training on anxiety: A meta-analysis and meta-regression analysis of randomized controlled trials. *Sports Med (Auckland)*. 2017;47(12):2521–32.
166. Cunha PM, Werneck AO, Santos LD, et al. Can resistance training improve mental health outcomes in older adults? A systematic review and meta-analysis of randomized controlled trials. *Psychiatry Res*. 2024;333:115746.
167. Gordon BR, McDowell CP, Hallgren M, Meyer JD, Lyons M, Herring MP. Association of efficacy of resistance exercise training with depressive symptoms: Meta-analysis and meta-regression analysis of randomized clinical trials. *JAMA Psychiatry*. 2018;75(6):566–76.
168. Strickland JC, Smith MA. The anxiolytic effects of resistance exercise. *Front Psychol*. 2014;5(753):753.
169. Gordon BR, McDowell CP, Lyons M, Herring MP. Resistance exercise training for anxiety and worry symptoms among young adults: A randomized controlled trial. *Sci Rep*. 2020;10(1):17548.
170. O'Sullivan D, Gordon BR, Lyons M, Meyer JD, Herring MP. Effects of resistance exercise training on depressive symptoms among young adults: A randomized controlled trial. *Psychiatry Res*. 2023;326:115322.
171. Cabanas-Sánchez V, Esteban-Cornejo I, Parra-Soto S, et al. Muscle strength and incidence of depression and anxiety: Findings from the UK Biobank prospective cohort study. *J Cachexia Sarcopenia Muscle*. 2022;13(4):1983–94.
172. Celis-Morales CA, Lyall DM, Anderson J, et al. The association between physical activity and risk of mortality is modulated by grip strength and cardiorespiratory fitness: Evidence from 498 135 UK-Biobank participants. *Eur Heart J*. 2017;38(2):116–22.
173. Ferrer-Ramos P, Garnacho-Castaño MV, Girabent-Farrés M, Faundez-Zanuy M, Serra-Payá N. Physical performance tests for preliminary cognitive screening in older adults: A systematic review of strength, walking, and balance assessments. *Arch Gerontol Geriatr*. 2025;130:105722.
174. Wang Y, Mu D, Wang Y. Association of low muscle mass with cognitive function and mortality in USA seniors: Results from NHANES 1999-2002. *BMC Geriatr*. 2024;24(1):420.
175. Hong AR, Kim SW. Effects of resistance exercise on bone health. *Endocrinol Metab (Seoul)*. 2018;33(4):435–44.
176. Souza D, Barbalho M, Ramirez-Campillo R, Martins W, Gentil P. High and low-load resistance training produce similar effects on bone mineral density of middle-aged and older people: A systematic review with meta-analysis of randomized clinical trials. *Exp Gerontol [Internet]*. 2020;138:110973.
177. Fritz N, Gene-Morales J, Saez-Berlanga A, Juesas A, Flández J, Colado JC. Resistance training for chronic low back pain in the elderly: A systematic review. *J Human Sport Exerc [Internet]*. 2021;16(3proc):S1492.
178. Vincent HK, George SZ, Seay AN, Vincent KR, Hurley RW. Resistance exercise, disability, and pain catastrophizing in obese adults with back pain. *Med Sci Sports Exerc*. 2014;46(9):1693–701.
179. Wasser JG, Vasilopoulos T, Zdziarski LA, Vincent HK. Exercise benefits for chronic low back pain in overweight and obese individuals. *PM R*. 2017;9(2):181–92.
180. Lim J, Choi A, Kim B. The effects of resistance training on pain, strength, and function in osteoarthritis. *J Pers Med*. 2024;14(12):1130.
181. Shiroma EJ, Cook NR, Manson JE, et al. Strength training and the risk of type 2 diabetes and cardiovascular disease. *Med Sci Sports Exerc*. 2017;49(1):40–6.

182. Tarp J, Støle AP, Blond K, Grøntved A. Cardiorespiratory fitness, muscular strength and risk of type 2 diabetes: A systematic review and meta-analysis. *Diabetologia*. 2019;62(7):1129–42.
183. Wei J, Fan L, Xia F, Zhu X, Chen L, Wang T. Association of aerobic and muscle-strengthening physical activity with all-cause and cardiovascular disease mortality among adults with type 2 diabetes: A prospective cohort of US adults. *Diabetes Metab Syndr*. 2024;18(3):102975.
184. Qadir R, Sculthorpe NF, Todd T, Brown EC. Effectiveness of resistance training and associated program characteristics in patients at risk for type 2 diabetes: A systematic review and meta-analysis. *Sports Med Open*. 2021;7(1):38.
185. Jansson AK, Chan LX, Lubans DR, Duncan MJ, Plotnikoff RC. Effect of resistance training on HbA1c in adults with type 2 diabetes mellitus and the moderating effect of changes in muscular strength: A systematic review and meta-analysis. *BMJ Open Diabetes Res Care*. 2022;10(2):e002595.
186. Consitt LA, Dudley C, Saxena G. Impact of endurance and resistance training on skeletal muscle glucose metabolism in older adults. *Nutrients*. 2019;11(11):2636.
187. Niemann MJ, Tucker LA, Bailey BW, Davidson LE. Strength training and insulin resistance: The mediating role of body composition. *J Diabetes Res*. 2020;2020:7694825.
188. Xie Y, Zhao H, Zhao M, et al. Effects of resistance exercise on blood glucose level and pregnancy outcome in patients with gestational diabetes mellitus: A randomized controlled trial. *BMJ Open Diabetes Res Care*. 2022;10(2):e002622.
189. Yang YR, Wang RY, Lin KH, Chu MY, Chan RC. Task-oriented progressive resistance strength training improves muscle strength and functional performance in individuals with stroke. *Clin Rehabil*. 2006;20(10):860–70.
190. Dalgas U, Stenager E. Exercise and disease progression in multiple sclerosis: Can exercise slow down the progression of multiple sclerosis? *Ther Adv Neurol Disord*. 2012;5(2):81–95.
191. Gomes Neto M, Conceição CS, Oliveira Carvalho V, Brites C. Effects of combined aerobic and resistance exercise on exercise capacity, muscle strength and quality of life in HIV-infected patients: A systematic review and meta-analysis. *PLoS One*. 2015;10(9):e0138066.
192. Zanetti HR, Lopes LTP, Gonçalves A, et al. Effects of resistance training on muscle strength, body composition and immune-inflammatory markers in people living with HIV: A systematic review and Meta-analysis of randomized controlled trials. *HIV Res Clin Pract*. 2021;22(5):119–27.
193. Chen N, He X, Feng Y, Ainsworth BE, Liu Y. Effects of resistance training in healthy older people with sarcopenia: A systematic review and meta-analysis of randomized controlled trials. *Eur Rev Aging Phys Act*. 2021;18(1):23.
194. Jiménez-Lupión D, Chirosa-Ríos L, Martínez-García D, Rodríguez-Pérez M, Jerez-Mayorga D. Effects of power training on functional capacity related to fall risk in older adults: A systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2023;104(9):1514–25.
195. Kaufman M, Dyrek P, Fredericson M, et al. The role of physical exercise in cognitive preservation: A systematic review. *Am J Lifestyle Med*. 2023;18(4):574–91.
196. Paluch AE, Boyer WR, Franklin BA, et al; on behalf the American Heart Association Council on Lifestyle and Cardiometabolic Health; Council on Arteriosclerosis, Thrombosis and Vascular Biology; Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; and Council on Peripheral Vascular Disease. Resistance exercise training in individuals with and without cardiovascular disease: 2023 update: A scientific statement from the American heart association. *Circulation*. 2024;149(3):e217–31.
197. Pillay J, Gaudet LA, Saba S, et al. Falls prevention interventions for community-dwelling older adults: Systematic review and meta-analysis of benefits, harms, and patient values and preferences. *Syst Rev* 2024;13(1):289.
198. Sauvage LR, Myklebust BM, Crow-Pan J, et al. A clinical trial of strengthening and aerobic exercise to improve gait and balance in elderly male nursing home residents. *Am J Phys Med Rehabil*. 1992;71(6):333–42.
199. Skelton DA, McLaughlin AW. Training functional ability in old age. *Physiother3* 1996;82(3):159–67.
200. Yoon DH, Lee J-Y, Song W. Effects of resistance exercise training on cognitive function and physical performance in cognitive frailty: A randomized controlled trial. *J Nutr Health Aging*. 2018;22(8):944–51.
201. Grgic J, Lazinica B, Schoenfeld BJ, Pedisic Z. Test-retest reliability of the one-repetition maximum (1RM) strength assessment: A systematic review. *Sports Med Open*. 2020;6(1):31.
202. Warneke K, Lohmann LH, Lima CD, et al. Physiology of stretch-mediated hypertrophy and strength increases: A narrative review. *Sports Med*. 2023;53(11):2055–75.
203. Abdul-Hameed U, Rangra P, Shareef MY, Hussain ME. Reliability of 1-repetition maximum estimation for upper and lower body muscular strength measurement in untrained middle aged type 2 diabetic patients. *Asian J Sports Med*. 2012;3(4):267–73.
204. Beia R, Wassermann A, Raps S, Mayhew J, Uder M, Kemmler W. Developing accurate repetition prediction equations for trained older adults with osteopenia. *Sports (Basel)*. 2024;12(9):233.
205. Fashi M, Shah Hosseini J, Ahmadi S. Predicting one repetition maximum in novice males: An RPE-based bench press model. *J Bodyw Mov Ther*. 2024;40:959–64.
206. Kemmler WK, Lauber D, Wassermann A, Mayhew JL. Predicting maximal strength in trained postmenopausal woman. *J Strength Cond Res*. 2006;20(4):838–42.
207. LeMense AT, Malone GT, Kinderman MA, Fedewa MV, Winchester LJ. Validity of using the load-velocity relationship to estimate 1 repetition maximum in the back squat exercise: A systematic review and meta-analysis. *J Strength Cond Res*. 2024;38(3):612–9.
208. Ramos AG. Resistance training intensity prescription methods based on lifting velocity monitoring. *Int J Sports Med*. 2023;45(4):257–66.
209. Kim PS, Mayhew JL, Peterson DF. A modified YMCA bench press test as a predictor of 1 repetition maximum bench press strength. *J Strength Cond Res*. 2002;16(3):440–5.
210. Golding L. *YMCA Fitness Testing and Assessment Manual*. 4th ed. Human Kinetics; 2000.
211. Ronai P. The YMCA bench press test. *ACSM'S Health & Fitness Journal*. 2020;24(6):33–6.
212. Gismondi A, Iellamo F, Caminiti G, et al. Rate of perceived exertion based on repetitions in reserve versus percentage of one-repetition maximum for resistance training prescription in cardiac rehabilitation: A pilot study. *J Cardiovasc Dev Dis*. 2024;12(1):8.
213. Elsangedy HM, Krinski K, Machado DG, Agricola PM, Okano AH, Gregorio da Silva S. Self-selected intensity, ratings of perceived exertion, and affective responses in sedentary male subjects during resistance training. *J Phys Ther Sci*. 2016;28(6):1795–800.
214. Warneke K, Wagner C-M, Keiner M, et al. Maximal strength measurement: A critical evaluation of common methods—a narrative review. *Front Sports Act Living*. 2023;5:1105201.
215. Jackson SM, Cheng MS, AR S Jr, Kolber MJ. Intrarater reliability of hand held dynamometry in measuring lower extremity isometric strength using a portable stabilization device. *Musculoskelet Sci Pract*. 2017;27:137–41.

216. McGrath R, Cawthon PM, Clark BC, Fielding RA, Lang JJ, Tomkinson GR. Recommendations for reducing heterogeneity in handgrip strength protocols. *J Frailty Aging*. 2022;11(2):143–50.
217. Mentiplay BF, Perraton LG, Bower KJ, et al. Assessment of lower limb muscle strength and power using hand-held and fixed dynamometry: A reliability and validity study. *PLoS One*. 2015;10(10):e0140822.
218. Quattrocchi A, Garufi G, Gugliandolo G, et al. Handgrip strength in health applications: A review of the measurement methodologies and influencing factors. *Sensors*. 2024;24(16):5100.
219. Schaubert KL, Bohannon RW. Reliability and validity of three strength measures obtained from community-dwelling elderly persons. *J Strength Cond Res*. 2005;19(3):717–20.
220. Bohannon RWR, Steffl M, Glenney S, et al. The prone bridge test: Performance, validity, and reliability among older and younger adults. *J Bodyw Mov Ther*. 2018;22(2):385. doi:10.1016/j.jbmt.2017.07.005
221. Behm DG, Daneshjoo A, Alizadeh S. Assessments of core fitness. *ACSM's Health Fit. J*. 2022;26(5):68–83.
222. Gibson A, Wagner D, Heyward V. *Advanced Fitness Assessment and Exercise Prescription*. 8th ed. Human Kinetics Publishers Inc; 2019.
223. Rikli RE. Jones CJ development and validation of a functional fitness test for community-residing older adults. *JAPA*. 1999;7:129–61.
224. Rikli RE, Jones CJ. Development and validation of criterion-referenced clinically relevant fitness standards for maintaining physical independence in later years. *Gerontologist*. 2013;53(2):255–67.
225. Rikli RE, Jones CJ. *Senior Fitness Test Manual*. Human Kinetics; 2013.
226. Lee SY, Gallagher D. Assessment methods in human body composition. *Curr Opin Clin Nutr Metab Care*. 2008;11(5):566–72.
227. Lohman TG, Milliken LA. *ACSM's Body Composition Assessment*. Human Kinetics; 2020.
228. Lim LS, Hoeksema LJ, Sherin K; ACPM Prevention Practice Committee. Screening for osteoporosis in the adult U.S. population: ACPM position statement on preventive practice. *Am J Prev Med* 2009;36(4):366–75.
229. Wally SF, Wali MF, Hariri OA, et al. Comparative diagnostic performance of dual-energy X-ray absorptiometry and other radiological modalities in osteoporosis detection: A systematic review. *J Clin Densitom*. 2025;29(1):101637.
230. Deshpande N, Hadi MS, Lillard JC, et al. Alternatives to DEXA for the assessment of bone density: A systematic review of the literature and future recommendations. *J Neurosurg Spine*. 2023;38(4):436–45.
231. Dhainaut A, Hoff M, Syversen U, Haugeberg G. Technologies for assessment of bone reflecting bone strength and bone mineral density in elderly women: An update. *Womens Health (Lond)*. 2016;12(2):209–16.
232. Bellicha A, van Baak MA, Battista F, et al. Effect of exercise training on weight loss, body composition changes, and weight maintenance in adults with overweight or obesity: An overview of 12 systematic reviews and 149 studies. *Obes Rev*. 2021;22(Suppl 4):e13256.
233. Armstrong A, Jungbluth Rodriguez K, Sabag A, et al. Effect of aerobic exercise on waist circumference in adults with overweight or obesity: A systematic review and meta-analysis. *Obes Rev*. 2022;23(8):e13446.
234. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK; American College of Sports Medicine. American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc*. 2009;41(2):459–71.
235. Jakicic JM, Powell KE, Campbell WW, et al; 2018 PHYSICAL ACTIVITY GUIDELINES ADVISORY COMMITTEE*. Physical activity and the prevention of weight gain in adults: A systematic review. *Med Sci Sports Exerc*. 2019;51(6):1262–9.
236. Jensen MD, Ryan DH, Apovian CM, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. AHA/ACC/TOS guideline for the management of overweight and obesity in adults. *Circulation*. 20132014;129(25 Suppl 2):S102–38.
237. Jakicic JM, Apovian CM, Barr-Anderson DJ, et al. Physical activity and excess body weight and adiposity for adults. American college of sports medicine consensus statement. *Med Sci Sports Exerc*. 2024;56(10):2076–91.
238. Vissers D, Hens W, Taeymans J, Baeyens JP, Poortmans J, Gaal L. The effect of exercise on visceral adipose tissue in overweight adults: A systematic review and meta-analysis. *PLoS One*. 2013;8(2):e56415.
239. Wewege MA, Desai I, Honey C, et al. The effect of resistance training in healthy adults on body fat percentage, fat mass and visceral fat: A systematic review and meta-analysis. *Sports Med*. 2022;52(2):287–300.
240. Jayedi A, Soltani S, Motlagh SZ-T, et al. Anthropometric and adiposity indicators and risk of type 2 diabetes: Systematic review and dose-response meta-analysis of cohort studies. *BMJ*. 2022;376:e067516.
241. Neeland IJ, Ross R, Després J-P, et al; International Atherosclerosis Society. Visceral and ectopic fat, atherosclerosis, and cardiometabolic disease: A position statement. *Lancet Diabetes Endocrinol*. 2019;7(9):715–25.
242. Escriche-Escuder A, Fuentes-Abolafio IJ, Roldán-Jiménez C, Cuesta-Vargas AI. Effects of exercise on muscle mass, strength, and physical performance in older adults with sarcopenia: A systematic review and meta-analysis according to the EWGSOP criteria. *Exp Gerontol*. 2021;151:111420.
243. Chaves TS, Scarpelli MC, Bergamasco JGA, et al. Effects of resistance training overload progression protocols on strength and muscle mass. *Int J Sports Med*. 2024;45(7):504–10.
244. Marzuca-Nassr GN, Alegría-Molina A, SanMartín-Calísto Y, et al. Muscle mass and strength gains following resistance exercise training in older adults 65-75 years and older adults above 85 years. *Int J Sport Nutr Exerc Metab*. 2023;34(1):11–9.
245. Viecelli C, Ewald CY. The non-modifiable factors age, gender, and genetics influence resistance exercise. *Front Aging*. 2022;3:1005848.
246. Al-Mhanna SB, Batrakoulis A, Norhayati MN, et al. Combined aerobic and resistance training improves body composition, alters cardiometabolic risk, and ameliorates cancer-related indicators in breast cancer patients and survivors with overweight/obesity: A systematic review and meta-analysis of randomized controlled trials. *J Sports Sci Med*. 2024;23(2):366–95.
247. Davidson LE, Hudson R, Kilpatrick K, et al. Effects of exercise modality on insulin resistance and functional limitation in older adults: A randomized controlled trial. *Arch Intern Med*. 2009;169(2):122–31.
248. Mathisen TF, Ackland T, Burke LM, et al. Best practice recommendations for body composition considerations in sport to reduce health and performance risks: A critical review, original survey and expert opinion by a subgroup of the IOC consensus on relative energy deficiency in sport (REDs. *Br J Sports Med*. 2023;57(17):1148–58.
249. Popp CJ, Jesch ED. The relationship between cardiorespiratory fitness and indices of fat mass and fat-free mass in adults. *Front Sports Act Living*. 2025;7:1583432.

250. Mondal H, Mishra SP. Effect of BMI, body fat percentage and fat free mass on maximal oxygen consumption in healthy young adults. *J Clin Diagn Res*. 2017;11(6):CC17–20.
251. Riebe D, Blissmer BJ, Greaney ML, Garber CE, Lees FD, Clark PG. The relationship between obesity, physical activity, and physical function in older adults. *J Aging Health*. 2009;21(8):1159–78.
252. Avgerinos KI, Spyrou N, Mantzoros CS, Dalamaga M. Obesity and cancer risk: Emerging biological mechanisms and perspectives. *Metabolism*. 2019;92:121–35.
253. Casanova F, O'Loughlin J, Martin S, et al. Higher adiposity and mental health: Causal inference using Mendelian randomization. *Hum Mol Genet*. 2021;30(24):2371–82.
254. Dwivedi AK, Dubey P, Cistola DP, Reddy SY. Association between obesity and cardiovascular outcomes: Updated evidence from meta-analysis studies. *Curr Cardiol Rep*. 2020;22(4):25.
255. Heymsfield SB, Wadden TA. Mechanisms, pathophysiology, and management of obesity. *N Engl J Med*. 2017;376(3):254–66.
256. Kim MS, Kim WJ, Khera AV, et al. Association between adiposity and cardiovascular outcomes: An umbrella review and meta-analysis of observational and Mendelian randomization studies. *Eur Heart J*. 2021;42(34):3388–403.
257. Powell-Wiley TM, Baumer Y, Baah FO, et al. Social determinants of cardiovascular disease. *Circ Res*. 2022;130(5):782–99.
258. Pulit SL, Stoneman C, Morris AP, et al. Meta-analysis of genome-wide association studies for body fat distribution in 694 649 individuals of European ancestry. *Hum Mol Genet*. 2019;28(1):166–74.
259. Rost S, Freuer D, Peters A, et al. New indexes of body fat distribution and sex-specific risk of total and cause-specific mortality: A prospective cohort study. *BMC Public Health* 2018;18(1):427.
260. Britton KA, Massaro JM, Murabito JM, Kreger BE, Hoffmann U, Fox CS. Body fat distribution, incident cardiovascular disease, cancer, and all-cause mortality. *J Am Coll Cardiol*. 2013;62(10):921–5.
261. Correa-de-Araujo R, Addison O, Miljkovic I, et al. Myosteatosis in the context of skeletal muscle function deficit: An interdisciplinary workshop at the national institute on aging. *Front Physiol*. 2020;11:963.
262. Du X, Hidayat K, Shi B-M. Abdominal obesity and gastroesophageal cancer risk: Systematic review and meta-analysis of prospective studies. *Biosci Rep*. 2017;37(3):BSR20160474.
263. Du Y-Z, Yang J-Q, Tang J, Zhang C-T, Liu Y-F. Association between the skeletal muscle-to-visceral fat ratio and kidney stones: A cross-sectional study. *Front Nutr*. 2025;12:1549047.
264. Kyrgiou M, Kalliala I, Markozannes G, et al. Adiposity and cancer at major anatomical sites: Umbrella review of the literature. *BMJ*. 2017;356:j477.
265. Oguntade AS, Islam N, Malouf R, et al. Body composition and risk of incident heart failure in 1 million adults: A systematic review and dose-response meta-analysis of prospective cohort studies. *J Am Heart Assoc*. 2023;12(13):e029062.
266. Oguntade AS, Taylor H, Lacey B, Lewington S. Adiposity, fat-free mass and incident heart failure in 500 000 individuals. *Open Heart*. 2024;11(2):e002711.
267. Shungin D, Winkler TW, Croteau-Chonka DC, et al; ADIPOGen Consortium. New genetic loci link adipose and insulin biology to body fat distribution. *Nature*. 2015;518(7538):187–96.
268. Santana F, MO P, NY T, RMR P. Low muscle mass in older adults and mortality: A systematic review and meta-analysis. *Exp Gerontol*. 2021;152:111461.
269. Zhou HH, Liao Y, Peng Z, Liu F, Wang Q, Yang W. Association of muscle wasting with mortality risk among adults: A systematic review and meta-analysis of prospective studies. *J Cachexia Sarcopenia Muscle*. 2023;14(4):1596–612.
270. Camargo Pereira C, Pagotto V, de Oliveira C, Silveira EA. Low muscle mass and mortality risk later in life: A 10-year follow-up study. *PLoS One*. 2022;17(7):e0271579.
271. Carmeli E, Reznick AZ, Coleman R, Carmeli V. Muscle strength and mass of lower extremities in relation to functional abilities in elderly adults. *Gerontology*. 2000;46(5):249–57.
272. Kirk B, Harrison SL, Zanker J, et al. Interactions between bone density and muscle mass in predicting all-cause mortality: A 10-year prospective cohort study of 1388 older men (aged 77-101 years). *Age Ageing*. 2025;54(7):afaf189.
273. Li R, Xia J, Zhang XI, et al. Associations of muscle mass and strength with all-cause mortality among US older adults. *Med Sci Sports Exerc*. 2018;50(3):458–67.
274. Baumert P, Mäntyselkä S, Schönfelder M, et al. Skeletal muscle hypertrophy rewires glucose metabolism: an experimental investigation and systematic review. *J Cachexia Sarcopenia Muscle*. 2024;15(3):989–1002.
275. Frampton J, Murphy KG, Frost G, Chambers ES. Short-chain fatty acids as potential regulators of skeletal muscle metabolism and function. *Nat Metab*. 2020;2(9):840–8.
276. Fritzen AM, Lundsgaard A-M, Kiens B. Tuning fatty acid oxidation in skeletal muscle with dietary fat and exercise. *Nat Rev Endocrinol*. 2020;16(12):683–96.
277. Brown JC, Harhay MO, Harhay MN. Sarcopenia and mortality among a population-based sample of community-dwelling older adults. *J Cachexia Sarcopenia Muscle*. 2016;7(3):290–8.
278. Asano Y, Tsunoda K, Nagata K, et al. Association of muscle mass, muscle quality, and function with mortality in community-dwelling older adults: Focus on segmental phase angle and extracellular to intracellular water ratio. *Clin Nutr ESPEN*. 2025;70:240–8.
279. Vendrami C, Gonzalez Rodriguez E, Gatineau G, et al. Prediction of 10-year fragility fractures using muscle health indicators in postmenopausal women: The osteolaus cohort. *J Cachexia Sarcopenia Muscle*. 2025;16(3):e13837.
280. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Writing group for the european working group on sarcopenia in older people 2 (EWGSOP2), and the extended group for EWGSOP2. *Sarcopenia*. 2019;48(4):601.
281. Izquierdo M, Merchant RA, Morley JE, et al. International exercise recommendations in older adults (ICFSR): Expert consensus guidelines. *J Nutr Health Aging*. 2021;25(7):824–53.
282. Bosy-Westphal A, Müller MJ. Diagnosis of obesity based on body composition-associated health risks-time for a change in paradigm. *Obes Rev*. 2021;22(Suppl 2):e13190.
283. Busetto L, Dicker D, Frühbeck G, et al. A new framework for the diagnosis, staging and management of obesity in adults. *Nat Med*. 2024;30(9):2395–9.
284. Rubino F, Cummings DE, Eckel RH, et al. Definition and diagnostic criteria of clinical obesity. *Lancet Diabetes Endocrinol*. 2025;13(3):221–62.
285. Campa F, Toselli S, Mazzilli M, Gobbo LA, Coratella G. Assessment of body composition in athletes: A narrative review of available methods with special reference to quantitative and qualitative bioimpedance analysis. *Nutrients*. 2021;13(5):1620.
286. Lemos T, Gallagher D. Current body composition measurement techniques. *Curr Opin Endocrinol Diabetes Obes*. 2017;24(5):310–4.

287. Al-Gindan YY, Hankey CR, Leslie W, Govan L, Lean MEJ. Predicting muscle mass from anthropometry using magnetic resonance imaging as reference: A systematic review. *Nutr Rev*. 2014;72(2):113–26.
288. Heymsfield SB, Gonzalez MC, Lu J, Jia G, Zheng J. Skeletal muscle mass and quality: Evolution of modern measurement concepts in the context of sarcopenia. *Proc Nutr Soc*. 2015;74(4):355–66.
289. Walowski CO, Braun W, Maisch MJ, et al. Reference values for skeletal muscle mass—current concepts and methodological considerations. *Nutrients*. 2020;12(3):755.
290. Lee RC, Wang Z, Heo M, Ross R, Janssen I, Heymsfield SB. Total-body skeletal muscle mass: Development and cross-validation of anthropometric prediction models. *Am J Clin Nutr*. 2000;72(3):796–803.
291. Donath L, Faude O, Hagmann S, Roth R, Zahner L. Fundamental movement skills in preschoolers: A randomized controlled trial targeting object control proficiency. *Child Care Health Dev*. 2015;41(6):1179–87.
292. Logan SW, Ross SM, Chee K, Stodden DF, Robinson LE. Fundamental motor skills: A systematic review of terminology. *J Sports Sci*. 2018;36(7):781–96.
293. Bruton M, O'Dwyer N. Synergies in coordination: A comprehensive overview of neural, computational, and behavioral approaches. *J Neurophysiol*. 2018;120(6):2761–74.
294. Diedrichsen J, Shadmehr R, Ivry RB. The coordination of movement: Optimal feedback control and beyond. *Trends Cogn Sci*. 2010;14(1):31–9.
295. Kimura A, Yokozawa T, Ozaki H. Clarifying the biomechanical concept of coordination through comparison with coordination in motor control. *Front Sports Act Living*. 2021;3(753062):753062.
296. Delmas S, Casamento-Moran A, Park SH, Yacoubi B, Christou EA. Motor planning perturbation: Muscle activation and reaction time. *J Neurophysiol*. 2018;120(4):2059–65.
297. Sutter K, Oostwoud Wijdenes L, van Beers RJ, Medendorp WP. Movement preparation time determines movement variability. *J Neurophysiol*. 2021;125(6):2375–83.
298. Beamish D, Bhatti SA, MacKenzie IS, Wu J. Fifty years later: A neurodynamic explanation of Fitts' law. *J R Soc Interface*. 2006;3(10):649–54.
299. Marcori AJ, Monteiro PHM, Oliveira JA, Doumas M, Teixeira LA. Single leg balance training: A systematic review. *Percept Mot Skills*. 2022;129(2):232–52.
300. McLaughlin EC, El-Kotob R, Chaput J-P, et al. Balance and functional training and health in adults: An overview of systematic reviews. *Appl Physiol Nutr Metab*. 2020;45(10 (Suppl. 2)):S180–96.
301. Thomas E, Battaglia G, Patti A, et al. Physical activity programs for balance and fall prevention in elderly: A systematic review. *Medicine (Baltim)*. 2019;98(27):e16218.
302. Keating CJ, Cabrera-Linares JC, Párraga-Montilla JA, Latorre-Román PA, Del Castillo RM, García-Pinillos F. Influence of resistance training on gait & balance parameters in older adults: A systematic review. *Int J Environ Res Public Health*. 2021;18(4):1759.
303. Šarabon N, Kozinc Z. Effects of resistance exercise on balance ability: Systematic review and meta-analysis of randomized controlled trials. *Life (Basel)*. 2020;10(11):284.
304. Zhen K, Zhang S, Tao X, Li G, Lv Y, Yu L. A systematic review and meta-analysis on effects of aerobic exercise in people with Parkinson's disease. *NPJ Parkinsons Dis*. 2022;8(1):146.
305. Hwang PW-N, Braun KL. The effectiveness of dance interventions to improve older adults' health: A systematic literature review. *Altern Ther Health Med*. 2015;21(5):64–70.
306. Lu J, Abd Rahman NA, Wyon M, Shaharudin S. The effects of dance interventions on physical function and quality of life among middle-aged and older adults: A systematic review. *PLoS One*. 2024;19(4):e0301236.
307. Lesinski M, Hortobágyi T, Muehlbauer T, Gollhofer A, Granacher U. Effects of balance training on balance performance in healthy older adults: A systematic review and meta-analysis. *Sports Med*. 2015;45(12):1721–38.
308. Lesinski M, Hortobágyi T, Muehlbauer T, Gollhofer A, Granacher U. Dose-response relationships of balance training in healthy young adults: A systematic review and meta-analysis. *Sports Med (Auckland)* 2015;45(4):557–76.
309. Castillo de Lima V, Castaño LAA, Boas VV, Uchida MC. A training program using an agility ladder for community-dwelling older adults. *J Vis Exp*. 2020;(157). doi:10.3791/60468
310. Lichtenstein E, Morat M, Roth R, Donath L, Faude O. Agility-based exercise training compared to traditional strength and balance training in older adults: A pilot randomized trial. *PeerJ*. 2020;8:e8781.
311. Lichtenstein E, Held S, Rappelt L, et al. Agility training to integratively promote neuromuscular, cardiorespiratory and cognitive function in healthy older adults: A one-year randomized-controlled trial. *Eur Rev Aging Phys Act*. 2023;20(1):21.
312. Morat M, Morat T, Zijlstra W, Donath L. Effects of multimodal agility-like exercise training compared to inactive controls and alternative training on physical performance in older adults: A systematic review and meta-analysis. *Eur Rev Aging Phys Act* 2021;18(1):4.
313. Castillo de Lima V, Castaño LAA, Sampaio RAC, Sampaio PYS, Teixeira CVL, Uchida MC. Effect of agility ladder training with a cognitive task (dual task) on physical and cognitive functions: A randomized study. *Front Public Health*. 2023;11:1159343.
314. Abellan van Kan G, Rolland Y, Andrieu S, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people an international academy on nutrition and aging (IANA) task force. *J Nutr Health Aging*. 2009;13(10):881–9.
315. Gallardo-Gómez D, Del Pozo-Cruz J, Noetel M, Álvarez-Barbosa F, Alfonso-Rosa RM, Del Pozo Cruz B. Optimal dose and type of exercise to improve cognitive function in older adults: A systematic review and bayesian model-based network meta-analysis of RCTs. *Ageing Res Rev*. 2022;76:101591.
316. Kamiya K, Hamazaki N, Matsue Y, et al. Gait speed has comparable prognostic capability to six-minute walk distance in older patients with cardiovascular disease. *Eur J Prev Cardiol*. 2018;25(2):212–9.
317. Lanzi S, Boichat J, Calanca L, Aubertin P, Malatesta D, Mazzolai L. Gait changes after supervised exercise training in patients with symptomatic lower extremity peripheral artery disease. *Vasc Med*. 2021;26(3):259–66.
318. Lyu X, Li S, Peng S, Cai H, Liu G, Ran X. Intensive walking exercise for lower extremity peripheral arterial disease: A systematic review and meta-analysis. *J Diabetes*. 2016;8(3):363–77.
319. Papa EV, Dong X, Hassan M. Resistance training for activity limitations in older adults with skeletal muscle function deficits: A systematic review. *Clin Interv Aging*. 2017;12:955–61.
320. Barker AL, Bird M-L, Talevski J. Effect of pilates exercise for improving balance in older adults: A systematic review with meta-analysis. *Arch Phys Med Rehabil*. 2015;96(4):715–23.
321. Behm DG, Aragão-Santos JC, Korooshfard N, Anvar SH. Alternative flexibility training. *Int J Sports Phys Ther* 2023;18(2):285–7.

322. Chen S, Zhang Y, Wang YT, Liu XL. Traditional chinese mind and body exercises for promoting balance ability of old adults: A systematic review and meta-analysis. *Evid Based Complement Alternat Med*. 2016;2016:7137362.
323. Chittrakul J, Siviroj P, Sungkarat S, Sapbamrer R. Multisystem physical exercise intervention for fall prevention and quality of life in pre-frail older adults: A randomized controlled trial. *Int J Environ Res Public Health*. 2020;17(9):3102.
324. Howe TE, Shea B, Dawson LJ, et al. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database Syst Rev*. 2011;2011(7):CD000333.
325. Howe TE, Rochester L, Neil F, Skelton DA, Ballinger C. Exercise for improving balance in older people. *Cochrane Database Syst Rev*. 2011;2011(11):CD004963.
326. Jadczak AD, Makwana N, Luscombe-Marsh N, Visvanathan R, Schultz TJ. Effectiveness of exercise interventions on physical function in community-dwelling frail older people: An umbrella review of systematic reviews. *JBI Database Syst Rev Implement Rep.* 2018;16(3):752–75.
327. Kan L, Zhang J, Yang Y, Wang P. The effects of Yoga on pain, mobility, and quality of life in patients with knee osteoarthritis: A systematic review. *Evid Based Complement Alternat Med*. 2016;2016:6016532.
328. Letton ME, Thom JM, Ward RE. The effectiveness of classical ballet training on health-related outcomes: A systematic review. *J Phys Act Health*. 2020;17(5):566–74.
329. Loewenthal J, Innes KE, Mitzner M, Mita C, Orkaby AR. Effect of yoga on frailty in older adults: A systematic review. *Ann Intern Med*. 2023;176(4):524–35.
330. Markovic G, Sarabon N, Greblo Z, Krizanic V. Effects of feedback-based balance and core resistance training vs. Pilates training on balance and muscle function in older women: A randomized-controlled trial. *Arch Gerontol Geriatr*. 2015;61(2):117–23.
331. Neves LM, Fortaleza AC, Rossi FE, et al. Functional training reduces body fat and improves functional fitness and cholesterol levels in postmenopausal women: A randomized clinical trial. *J Sports Med Phys Fitness*. 2017;57(4):448–56.
332. Shen Y, Shi Q, Nong K, et al. Exercise for sarcopenia in older people: A systematic review and network meta-analysis. *J Cachexia Sarcopenia Muscle*. 2023;14(3):1199–211.
333. Widjaja W, Wongwattanapong T, Laskin JJ, Ajjimaporn A. Benefits of thai yoga on physical mobility and lower limb muscle strength in overweight/obese older women: A randomized controlled pilot trial. *Complement Ther Clin Pract*. 2021;43:101345.
334. Wolf R, Locks RR, Lopes PB, et al. Multicomponent exercise training improves gait ability of older women rather than strength training: A randomized controlled trial. *J Aging Res*. 2020;2020:6345753.
335. Yang Y, Schooten KS, Komisar V, et al. Effects of the mobility-fit physical activity program on strength and mobility in older adults in assisted living: A feasibility study. *Int J Environ Res Public Health*. 2022;19(9):5453.
336. Zemková E, Zapletalová L. The role of neuromuscular control of postural and core stability in functional movement and athlete performance. *Front Physiol*. 2022;13:796097.
337. Lee NG, Kang TW, Park HJ. Relationship between balance, gait, and activities of daily living in older adults with dementia. *Geriatr Orthop Surg Rehabil*. 2020;11:2151459320929578.
338. Olivares-Jabalera J, Fílter-Ruger A, Dos'Santos T, et al. Exercise-based training strategies to reduce the incidence or mitigate the risk factors of anterior cruciate ligament injury in adult football (soccer) players: A systematic review. *Int J Environ Res Public Health*. 2021;18(24):13351.
339. Wang Q, Jiang X, Shen Y, et al. Effectiveness of exercise intervention on fall-related fractures in older adults: A systematic review and meta-analysis of randomized controlled trials. *BMC Geriatr*. 2020;20(1):322.
340. Neptune R, Vistamehr A. Dynamic balance during human movement: Measurement and control mechanisms. *J Biomech Eng*. 2018;141(7):0708011–07080110.
341. Allen JL, Franz JR. The motor repertoire of older adult fallers may constrain their response to balance perturbations. *J Neurophysiol*. 2018;120(5):2368–78.
342. Xing L, Bao Y, Wang B, et al. Falls caused by balance disorders in the elderly with multiple systems involved: Pathogenic mechanisms and treatment strategies. *Front Neurol*. 2023;14:1128092.
343. Jiang J, Zhou Q, Zhang C, Cong K. Activities of daily living mediate the association between balance and falls in middle aged and older adults. *Sci Rep*. 2025;15(1):28694.
344. Kozinc Z, Löfler S, Hofer C, Carraro U, Šarabon N. Diagnostic balance tests for assessing risk of falls and distinguishing older adult fallers and non-fallers: A systematic review with meta-analysis. *Diagnostics (Basel)*. 2020;10(9):667.
345. Yang Y, Zeng Z, van Schooten KS, et al. Effects of a multicomponent physical activity programme, Mobility-Fit, compared with a standard care lower limb strengthening programme, to promote safe mobility among older adults in care facilities: Protocol for a cluster randomised controlled trial. *BMJ Open*. 2024;14(1):e082403.
346. Rodrigues F, Monteiro AM, Forte P, Morouço P. Effects of muscle strength, agility, and fear of falling on risk of falling in older adults. *Int J Environ Res Public Health*. 2023;20(6):4945.
347. Deshpande N, Metter EJ, Lauretani F, Bandinelli S, Ferrucci L. Interpreting fear of falling in the elderly: what do we need to consider? *J Geriatr Phys Ther*. 2009;32(3):91–6.
348. da Silva RG, da Silva Sobrinho AC, Costa GP, et al. Benefits of physical exercise through multivariate analysis in sedentary adults and elderly: An analysis of physical fitness, health and anthropometrics. *Exp Gerontol*. 2025;200:112669.
349. Hornyak V, Brach JS, Wert DM, Hile E, Studenski S, VanSwearingen JM. What is the relation between fear of falling and physical activity in older adults? *Arch Phys Med Rehabil*. 2013;94(12):2529–34.
350. Garcia Meneguci CA, Meneguci J, Sasaki JE, Tribess S, Júnior JSV. Physical activity, sedentary behavior and functionality in older adults: A cross-sectional path analysis. *PLoS One*. 2021;16(1):e0246275.
351. Chung P-K, Zhao Y, Liu J-D, Quach B. A canonical correlation analysis on the relationship between functional fitness and health-related quality of life in older adults. *Arch Gerontol Geriatr*. 2017;68:44–8.
352. Denfeld QE, Turrise S, MacLaughlin EJ, et al; on behalf of the American Heart Association Cardiovascular Disease in Older Populations Committee of the Council on Clinical Cardiology and Council on Cardiovascular and Stroke Nursing; Council on Lifestyle and Cardiometabolic Health; and Stroke Council. Preventing and managing falls in adults with cardiovascular disease: A scientific statement from the American heart association. *Circ Cardiovasc Qual Outcomes*. 2022;15(6):e000108.
353. Lee O, Park D. Walking speed and hand grip strength as predictors of hypertension risk in older adults: The impact of insulin resistance. *Geriatr Gerontol Int*. 2025;25(12):1923–9.
354. Ramnath U, Rauch L, Lambert EV, Kolbe-Alexander TL. The relationship between functional status, physical fitness and cognitive performance in physically active older adults: A pilot study. *PLoS One*. 2018;13(4):e0194918.

355. Wang Z, Cui K, Song R, et al. Influence of cardiovascular risk burden on motor function among older adults: Mediating role of cardiovascular diseases accumulation and cognitive decline. *Front Med (Lausanne)*. 2022;9:856260.
356. Yi Y, Park Y-H. Structural equation model of the relationship between functional ability, mental health, and quality of life in older adults living alone. *PLoS One*. 2022;17(8):e0269003.
357. Toth EE, Ihász F, Ruíz-Barquín R, Szabo A. Physical activity and psychological resilience in older adults: A systematic review of the literature. *J Aging Phys Act*. 2024;32(2):276–86.
358. Yang Q, Zhang Y, Li S. The impact of mind-body exercise on the quality of life in older adults: The chain mediation effect of perceived social support and psychological resilience. *Front Public Health*. 2024;12:1446295.
359. Donath L, van Dieën J, Faude O. Exercise-based fall prevention in the elderly: What about agility? *Sports Med*. 2016;46(2):143–9.
360. Kim MK, Kang BH, Park JH, Ham SM, Park HY, Hong I. Longitudinal changes in dual decline in memory and gait speed association with instrumental activities of daily living: Findings from the national social life, health, and aging project. *J Aging Health*. 2023;35(3-4):256–64.
361. Ma Lourdes DR-S, Sergio A-R, Francisco R-OJ, Blanco-Saez M. Gait speed assessment as a prognostic tool for morbidity and mortality in vulnerable older adult patients following vascular surgery. *Geriatr Nurs*. 2024;56:25–31.
362. Marín-Jiménez N, Cruz-León C, Perez-Bey A, et al. Predictive validity of motor fitness and flexibility tests in adults and older adults: A systematic review. *J Clin Med*. 2022;11(2):328.
363. McGinn AP, Kaplan RC, Verghese J, et al. Walking speed and risk of incident ischemic stroke among postmenopausal women. *Stroke*. 2008;39(4):1233–9.
364. McLaughlin EC, El-Kotob R, Chaput JP, et al. Balance and functional training and health in adults: An overview of systematic reviews. *Appl Physiol Nutr Metab*. 2020;45(10 Suppl. 2):S180–96.
365. Melzer I, Kurz I. Self reported function and disability in late life: A comparison between recurrent fallers and non-fallers. *Disabil Rehabil*. 2009;31(10):791–8.
366. Prommaban A, Moonkayaow S, Phinyo P, Siviroj P, Sirikul W, Lerttrakarnnon P. The effect of exercise program interventions on frailty, clinical outcomes, and biomarkers in older adults: A systematic review. *J Clin Med*. 2024;13(21):6570.
367. Luo Y, Hao J, Zhu L, et al. Effects of multicomponent exercise nursing intervention in elderly stroke patients with frailty: A randomized controlled trial. *Front Med (Lausanne)*. 2024;11:1450494.
368. Cordes T, Zwingmann K, Rudisch J, Voelcker-Rehage C, Wollesen B. Multicomponent exercise to improve motor functions, cognition and well-being for nursing home residents who are unable to walk—A randomized controlled trial. *Exp Gerontol*. 2021;153:111484.
369. Ibrahim A, Mat Ludin AF, Singh DKA, Rajab NF, Shahar S. Changes in cardiovascular-health blood biomarkers in response to exercise intervention among older adults with cognitive frailty: A scoping review. *Front Physiol*. 2023;14:1077078. doi:10.3389/fphys.2023.1077078
370. Figueira B, Bravo J, Raimundo A, Reis A. Effects of a 26 week multicomponent exercise program on cardiovascular and lipid profiles in premenopausal and postmenopausal women. *Sci Rep*. 2025;15(1):22485.
371. Savikangas T, Valadão P, Haapala EA, Finni T. Effects of multicomponent exercise intervention on cardiometabolic risk factors in children and young adults with cerebral palsy: A multiple-baseline trial. *BMC Sports Sci Med Rehabil*. 2024;16(1):219.
372. Subías-Perié J, Navarrete-Villanueva D, Fernández-García AI, et al. Effects of a multicomponent training followed by a detraining period on metabolic syndrome profile of older adults. *Exp Gerontol*. 2024;186:112363.
373. Voelcker-Rehage C, Niemann C. Structural and functional brain changes related to different types of physical activity across the life span. *Neurosci Biobehav Rev*. 2013;37(9 Pt B):2268–95.
374. Herold F, Hamacher D, Schega L, Müller NG. Thinking while moving or moving while thinking—concepts of motor-cognitive training for cognitive performance enhancement. *Front Aging Neurosci*. 2018;10:228.
375. Netz Y. Is there a preferred mode of exercise for cognition enhancement in older age?-a narrative review. *Front Med (Lausanne)*. 2019;6:57.
376. Arioli M, Tsantzalou KS, Qiu M, et al. The neural basis of sequence learning: A neuroimaging meta-analysis across social, cognitive and motor domains. *Neurosci Biobehav Rev*. 2026;181:106530.
377. Beck Jepsen D, Robinson K, Ogliari G, et al. Predicting falls in older adults: An umbrella review of instruments assessing gait, balance, and functional mobility. *BMC Geriatr*. 2022;22(1):615.
378. Colón-Emeric CS, McDermott CL, Lee DS, Berry SD. Risk assessment and prevention of falls in older community-dwelling adults: A review. *JAMA*. 2024;331(16):1397–406.
379. Loughran KJ, Atkinson G, Beauchamp MK, et al. Balance impairment in individuals with COPD: A systematic review with meta-analysis. *Thorax*. 2020;75(7):539–46.
380. Ozcan EB, Saglam M, Vardar-Yagli N, et al. Impaired balance and gait characteristics in patients with chronic heart failure. *Heart Lung Circ*. 2022;31(6):832–40.
381. de la Motte SJ, Lisman P, Gribbin TC, Murphy K, Deuster PA. Systematic review of the association between physical fitness and musculoskeletal injury risk: Part 3-flexibility, power, speed, balance, and agility. *J Strength Cond Res*. 2019;33(6):1723–35.
382. Figgins E, Choi Y-H, Speechley M, Montero-Odasso M. Associations between potentially modifiable and nonmodifiable risk factors and gait speed in middle- and older-aged adults: Results from the canadian longitudinal study on Aging. *J Gerontol A Biol Sci Med Sci*. 2021;76(10):e253–63.
383. Mehdipour A, Saunders S, Reid J, et al. Acceptability, reliability, and validity of virtually administered gait speed tests. *J Am Med Dir Assoc*. 2024;25(8):105048.
384. Sanders JB, Bremmer MA, Comijs HC, Ven PM, Deeg DJH, Beekman ATF. Gait speed and processing speed as clinical markers for geriatric health outcomes. *Am J Geriatr Psychiatry*. 2017;25(4):374–85.
385. Sato A, Kamiya K, Hamazaki N, et al. Gait speed and 6-minute walking distance are useful for identifying difficulties in activities of daily living in patients with cardiovascular disease. *Heart Lung*. 2022;51:46–51.
386. Önal B, Kocaman AA. The validity and reliability of the 10-meter walk test with obstacles in community-dwelling older adults. *Int J Rehabil Res*. 2025;48(2):120–5.
387. Berg KO, Wood-Dauphinee SL, Williams JI, Maki B. Measuring balance in the elderly: Validation of an instrument. *Can J Public Health*. 1992;83(Suppl 2):S7–11.
388. Duncan PW, Weiner DK, Chandler J, Studenski S. Functional reach: A new clinical measure of balance. *J Gerontol*. 1990;45(6):M192–197.
389. Phelan EA, Mahoney JE, Voit JC, Stevens JA. Assessment and management of fall risk in primary care settings. *Med Clin North Am*. 2015;99(2):281–93.
390. Plisky P, Schwartkopf-Phifer K, Huebner B, Garner MB, Bullock G. Systematic review and meta-analysis of the

Y-balance test lower quarter: Reliability, discriminant validity, and predictive validity. *Int J Sports Phys Ther*. 2021;16(5):1190–209.
391. Rose DJ, Lucchese N, Wiersma LD. Development of a multidimensional balance scale for use with functionally independent older adults. *Arch Phys Med Rehabil*. 2006;87(11):1478–85.
392. Young W, Rayner R, Talpey S. It's time to change direction on agility research: A call to action. *Sports Med Open*. 2021;7(1):12.
393. Bushman BA. Flexibility exercises and performance. *ACSMs Health Fit J*. 2016;20(5):5–9.
394. Elliott GF, Worthington CR. Muscle contraction: Viscous-like frictional forces and the impulsive model. *Int J Biol Macromol*. 2000;27(5):327–32.
395. Gleim GW, McHugh MP. Flexibility and its effects on sports injury and performance. *Sports Med (Auckland)*. 1997;24(5):289–99.
396. Konrad A, Alizadeh S, Daneshjoo A, et al. Chronic effects of stretching on range of motion with consideration of potential moderating variables: A systematic review with meta-analysis. *J Sport Health Sci*. 2024;13(2):186–94.
397. Konrad A, Nakamura M, Tilp M, Donti O, Behm DG. Foam rolling training effects on range of motion: A systematic review and meta-analysis. *Sports Med*. 2022;52(10):2523–35.
398. Alizadeh S, Daneshjoo A, Zahiri A, et al. Resistance training induces improvements in range of motion: A systematic review and meta-analysis. *Sports Med*. 2023;53(3):707–22.
399. Nikolaidis PT, Rosemann T, Knechtle B. Force-velocity characteristics, muscle strength, and flexibility in female recreational marathon runners. *Front Physiol*. 2018;9:1563.
400. Sell T, Tsai Y, Smoliga JM, Myers JB, Lephart SM. Strength, flexibility and balance characteristics of highly proficient golfers. *JSCR*. 2007;21(4):1166–71.
401. Behm DG, Kay AD, Trajano GS, Alizadeh S, Blazevich AJ. Effects of stretching on injury risk reduction and balance. 2021 [cited 2025 Apr 17]; Available from: https://jcep.kglmeridian.com/view/journals/ceph/10/3/article-p106.xml.
402. Behm DG, Blazevich AJ, Kay AD, McHugh M. Acute effects of muscle stretching on physical performance, range of motion, and injury incidence in healthy active individuals: A systematic review. *Appl Physiol Nutr Metab*. 2016;41(1):1–11.
403. Pfeifer CE, Ross LM, Weber SR, Sui X, Blair SN. Are flexibility and muscle-strengthening activities associated with functional limitation? *Sports Med Health Sci*. 2022;4(2):95–100.
404. Brito LBB, de Araújo DSMS, de Araújo CGS. Does flexibility influence the ability to sit and rise from the floor? *Am J Phys Med Rehabil*. 2013;92(3):241–7.
405. Hatefi M, Babakhani F, Ashrafizadeh M. The effect of static stretching exercises on hip range of motion, pain, and disability in patients with non-specific low back pain. *J Exp Orthop*. 2021;8(1):55.
406. Jeong Y, Heo S, Lee G, Park W. Pre-obesity and obesity impacts on passive joint range of motion. *Ergonomics*. 2018;61(9):1223–31.
407. Kell RT, Bell G, Quinney A. Musculoskeletal fitness, health outcomes and quality of life. *Sports Med*. 2001;31(12):863–73.
408. Misner JE, Massey BH, Bemben MG, Going S, Patrick J. Long-term effects of exercise on the range of motion of aging women. *J Orthop Sports Phys Ther*. 1992;16(1):37–42.
409. Park W, Ramachandran J, Weisman P, Jung ES. Obesity effect on male active joint range of motion. *Ergonomics*. 2010;53(1):102–8.
410. Wilke J, Macchi V, De Caro R, Stecco C. Fascia thickness, aging and flexibility: Is there an association? *J Anat*. 2019;234(1):43–9.
411. Steultjens MP, Dekker J, van Baar ME, Oostendorp RA, Bijlsma JW. Range of joint motion and disability in patients with osteoarthritis of the knee or hip. *Rheumatology (Oxford)*. 2000;39(9):955–61.
412. Oosterwijk AM, Nieuwenhuis MK, van der Schans CP, Mouton LJ. Shoulder and elbow range of motion for the performance of activities of daily living: A systematic review. *Physiother Theory Pract*. 2018;34(7):505–28.
413. Bible JE, Biswas D, Miller CP, Whang PG, Grauer JN. Normal functional range of motion of the lumbar spine during 15 activities of daily living. *J Spinal Disord Tech*. 2010;23(2):106–12.
414. Namdari S, Yagnik G, Ebaugh DD, et al. Defining functional shoulder range of motion for activities of daily living. *J Shoulder Elbow Surg*. 2012;21(9):1177–83.
415. Konrad A, Nakamura M, Sardroodian M, Aboozari N, Hadjizadeh Anvar S, Behm DG. The effects of chronic stretch training on musculoskeletal pain. *Eur J Appl Physiol [Internet]*. 2025;125(8):2037.
416. Arntz F, Markov A, Schoenfeld BJ, et al. Chronic effects of static stretching exercises on skeletal muscle hypertrophy in healthy individuals: A systematic review and multilevel meta-analysis. *Sports Med Open*. 2024;10(1):106.
417. Geneen LJ, Moore RA, Clark C, Martin D, Colvin LA, Smith BH. Physical activity and exercise for chronic pain in adults: An overview of Cochrane Reviews. *Cochrane Database Syst Rev* 2017;4(4):CD011279. [cited 2025 Mar 6]. Available from: https://pubmed.ncbi.nlm.nih.gov/28436583/. doi:10.1002/14651858.CD011279.pub3
418. Mertens MG, Meert L, Struyf F, Schwank A, Meeus M. Exercise therapy is effective for improvement in range of motion, function, and pain in patients with frozen shoulder: A systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2022;103(5):998–1012.e14.
419. Emilio EJM-L, Hita-Contreras F, Jiménez-Lara PM, Latorre-Román P, Martínez-Amat A. The association of flexibility, balance, and lumbar strength with balance ability: Risk of falls in older adults. *J Sports Sci Med*. 2014;13(2):349–57.
420. Takeuchi K, Nakamura M, Fukaya T, Nakao G, Mizuno T. Stretching intervention can prevent muscle injuries: A systematic review and meta-analysis. *Sport Sci Health*. 2024;20(4):1119–29.
421. Amako M, Oda T, Masuoka K, Yokoi H, Campisi P. Effect of static stretching on prevention of injuries for military recruits. *Mil Med*. 2003;168(6):442–6.
422. Hori M, Hasegawa H, Takasaki H. Comparisons of hamstring flexibility between individuals with and without low back pain: Systematic review with meta-analysis. *Physiother Theory Pract*. 2021;37(5):559–82.
423. Sadler SG, Spink MJ, Ho A, Jonge XJ, Chuter VH. Restriction in lateral bending range of motion, lumbar lordosis, and hamstring flexibility predicts the development of low back pain: A systematic review of prospective cohort studies. *BMC Musculoskelet Disord*. 2017;18(1):179.
424. Yamamoto K, Kawano H, Gando Y, et al. Poor trunk flexibility is associated with arterial stiffening. *Am J Physiol Heart Circ Physiol*. 2009;297(4):H1314–1318.
425. Bisconti AV, Cè E, Longo S, et al. Evidence for improved systemic and local vascular function after long-term passive static stretching training of the musculoskeletal system. *J Physiol*. 2020;598(17):3645–66.
426. Kato M, Masuda T, Ogano M, et al. Stretching exercises improve vascular endothelial dysfunction through attenuation of oxidative stress in chronic heart failure patients with an implantable cardioverter defibrillator. *J Cardiopulm Rehabil Prev*. 2017;37(2):130–8.

427. Thomas E, Bellafiore M, Gentile A, Paoli A, Palma A, Bianco A. Cardiovascular responses to muscle stretching: A systematic review and meta-analysis. *Int J Sports Med*. 2021;42(6):481.
428. Eda N, Ito H, Akama T. Beneficial effects of yoga stretching on salivary stress hormones and parasympathetic nerve activity. *J Sports Sci Med*. 2020;19(4):695–702.
429. Eda N, Ito H, Shimizu K, Suzuki S, Lee E, Akama T. Yoga stretching for improving salivary immune function and mental stress in middle-aged and older adults. *J Women Aging*. 2018;30(3):227–41.
430. Kruse NT, Scheuermann BW. Effect of self-administered stretching on NIRS-measured oxygenation dynamics. *Clin Physiol Funct Imaging*. 2016;36(2):126–33.
431. Matsuo H, Kubota M, Shimada S, et al. The effect of static stretching duration on muscle blood volume and oxygenation. *J Strength Cond Res*. 2022;36(2):379–85.
432. Wong A, Sanchez-Gonzalez M, Kalfon R, Alvarez-Alvarado S, Figueroa A. The effects of stretching training on cardiac autonomic function in obese postmenopausal women. *Altern Ther Health Med*. 2017;23(2):20–6.
433. Wong A, Figueroa A. Eight weeks of stretching training reduces aortic wave reflection magnitude and blood pressure in obese postmenopausal women. *J Hum Hypertens*. 2014;28(4):246–50.
434. Araújo CGS, de Souza E Silva CG, Kunutsor SK, et al. Reduced body flexibility is associated with poor survival in middle-aged men and women: A prospective cohort study. *Scand J Med Sci Sports*. 2024;34(8):e14708.
435. Hanks J, Myers B. Validity, reliability, and efficiency of a standard goniometer, medical inclinometer, and builder's inclinometer. *Int J Sports Phys Ther*. 2023;18(4):989–96.
436. Gandbhir VN, Cunha B. Goniometer. *StatPearls*. StatPearls Publishing; 2025 [cited 2026 Jan 10] Available from: http://www.ncbi.nlm.nih.gov/books/NBK558985/.
437. Soucie JM, Wang C, Forsyth A, et al; Hemophilia Treatment Center Network. Range of motion measurements: Reference values and a database for comparison studies. *Haemophilia*. 2011;17(3):500–7.
438. Hershkovich O, Grevitt MP, Lotan R. Schober test and its modifications revisited-what are we actually measuring? Computerized tomography-based analysis. *J Clin Med*. 2022;11(23):6895.
439. Jones CJ, Rikli RE, Max J, Noffal G. The reliability and validity of a chair sit-and-reach test as a measure of hamstring flexibility in older adults. *Res Q Exerc Sport*. 1998;69(4):338–43.
440. Lemmink KAPM, Kemper HCG, de Greef MHG, Rispens P, Stevens M. The validity of the sit-and-reach test and the modified sit-and-reach test in middle-aged to older men and women. *Res Q Exerc Sport*. 2003;74(3):331–6.
441. Mayorga-Vega D, Merino-Marban R, Viciana J. Criterion-related validity of sit-and-reach tests for estimating hamstring and lumbar extensibility: A meta-analysis. *J Sports Sci Med*. 2014;13(1):1–14.
442. Camino Willhuber GO, Piuzzi NS. Straight leg raise test. *StatPearls*. StatPearls Publishing; 2025 [cited 2026 Jan 11] Available from: http://www.ncbi.nlm.nih.gov/books/NBK539717/.
443. Powden CJ, Hoch JM, Hoch MC. Reliability and minimal detectable change of the weight-bearing lunge test: A systematic review. *Man Ther*. 2015;20(4):524–32.
444. Gajdosik RL, Bohannon RW. Clinical measurement of range of motion. Review of goniometry emphasizing reliability and validity. *Phys Ther*. 1987;67(12):1867–72.
445. Mullaney MJ, McHugh MP, Johnson CP, Tyler TF. Reliability of shoulder range of motion comparing a goniometer to a digital level. *Physiother Theory Pract*. 2010;26(5):327–33.
446. Jackson AW, Baker AA. The relationship of the sit and reach test to criterion measures of hamstring and back flexibility in young females. *Res Q Exerc Sport*. 1986;57(3):183–6.
447. Jackson A, Langford NJ. The criterion-related validity of the sit and reach test: Replication and extension of previous findings. *Res Q Exerc Sport*. 1989;60(4):384–7.
448. Muyor JM, Vaquero-Cristóbal R, Alacid F, López-Miñarro PA. Criterion-related validity of sit-and-reach and toe-touch tests as a measure of hamstring extensibility in athletes. *J Strength Cond Res*. 2014;28(2):546–55.
449. Gatton ML, Pearcy MJ. Kinematics and movement sequencing during flexion of the lumbar spine. *Clin Biomech (Bristol)*. 1999;14(6):376–83.
450. Alter MJ. *Science of Flexibility*. 3rd ed. Human Kinetics; 2004.
451. Sahrmann SA. *Diagnosis and Treatment of Movement Impairment Syndromes*. Mosby; 2002.
452. Warneke K, Lohmann LH, Behm DG, et al. Effects of chronic static stretching on maximal strength and muscle hypertrophy: A systematic review and meta-analysis with meta-regression. *Sports Med—Open*. 2024;10(1):45.
453. Ozaki H, Loenneke JP, Thiebaud R. Resistance training induced increase in VO2max in young and older subjects. *Eur Rev Aging Phys Act*. 2013;10:107–16.
454. Marques EA, Figueiredo P, Harris TB, Wanderley FA, Carvalho J. Are resistance and aerobic exercise training equally effective at improving knee muscle strength and balance in older women? *Arch Gerontol Geriatr*. 2017;68:106–12.
455. An J, Su Z, Meng S. Effect of aerobic training versus resistance training for improving cardiorespiratory fitness and body composition in middle-aged to older adults: A systematic review and meta-analysis of randomized controlled trials. *Arch Gerontol Geriatr*. 2024;126:105530.
456. Henderson RM, Leng XI, Chmelo EA, et al. Gait speed response to aerobic versus resistance exercise training in older adults. *Aging Clin Exp Res*. 2017;29(5):969–76.
457. Crane JD, Macneil LG, Tarnopolsky MA. Long-term aerobic exercise is associated with greater muscle strength throughout the life span. *J Gerontol A Biol Sci Med Sci*. 2013;68(6):631–8.
458. Vieira ND, Testa D, Ruas PC, Salvini T de F, Catai AM, de Melo RC. The effects of 12 weeks Pilates-inspired exercise training on functional performance in older women: A randomized clinical trial. *J Bodyw Mov Ther*. 2017;21(2):251–8.
459. Gerber M, Cheval B, Cody R, et al. Psychophysiological foundations of human physical activity behavior and motivation: Theories, systems, mechanisms, evolution, and genetics. *Physiol Rev*. 2025;105(3):1213–90.
460. Ekkekakis P. People have feelings! Exercise psychology in paradigmatic transition. *Curr Opin Psychol*. 2017;16:84–8.
461. Monahan K, Paiva A, Blissmer B, Sacco A, Robbins M. Transtheoretical model for exercise: Measure redevelopment and assessing the role of barriers in a diverse population. *Am J Health Promot*. 2024;38(2):186–96.
462. King AC, Powell KE, Kraus WE. The US physical activity guidelines advisory committee report-introduction. *Med Sci Sports Exerc*. 2019;51(6):1203–5.